

NET CONTENTS
5 LBS.

ZINEB-75 WP

PRODUCT
No. 978

| | |
|---|---------------|
| ACTIVE INGREDIENTS: | BY WT. |
| Zineb (Zinc ethylenebisdithiocarbamate) (Zinc expressed as metallic 17.7%) | 75.00% |
| INERT INGREDIENTS | 25.00% |
| TOTAL — 100.00% | |

® Trade Marks Reg. U. S. Patent Office

CAUTION: KEEP OUT OF THE REACH OF CHILDREN.

See below for additional cautions.

USDA Reg. No. 2342-716

FASCO ZINEB-75 WP is a wettable powder fungicidal product containing 75% ZINEB fungicide. It is designed for use in aqueous sprays against certain diseases attacking vegetable, fruit, and ornamental crops as described below.

DIRECTIONS

Use contrary to these directions may cause serious plant, crop or personal injury, unsatisfactory control, or other unintended consequences.

CITRUS (Oranges, Grapefruit, Tangerines) — To control Rust Mite and associated russeting and greasy spot, use 1 pound per 100 gallons of water. Apply with very thorough coverage. Make application from late June through early August or earlier if rust mite begins to appear.

TOMATOES — Against Early and Late Blight, Septoria leaf spot, Gray leaf spot (Stemphylium), Anthraco
se, use 1 1/2 to 2 pounds per 100 gallons of water. Apply when disease is first present and repeat ap-
plications at 7 to 10 day intervals or oftener if blight conditions are severe. Do not exceed 4 lbs. per
acre.

PEPPERS, EGGPLANT — Against Alternaria, Anthraco
se, and Phomopsis of eggplant; Anthraco
se and Cercospora leaf spot of pepper, use 1 1/2 to 2 pounds per 100 gallons of water. Start applications at first
sign of disease and repeat at 7 to 10 day intervals as needed. Do not exceed 4 lbs. per acre on Eggplant;
8 lbs. per acre on Pepper.

CABBAGE, BROCCOLI, CAULIFLOWER — Against Downy Mildew and Alternaria, use 1 1/2 to 2 pounds per
100 gallons of water. Apply at first sign of disease, repeat at 7 day intervals as needed. Do not apply
later than 7 days from harvest, do not exceed 6 lbs. per acre.

CANTALOUPE, CUCUMBERS, WATERMELONS, SQUASH — Against Downy Mildew, Anthraco
se, Gummy
Stem Blight, use 1 1/2 to 2 pounds per 100 gallons of water. Apply at first sign of disease and repeat at 7
to 10 day intervals as long as necessary. Under severe disease conditions, shorten interval to 3 to 5 days.
Do not exceed 4 lbs. per acre.

TOBACCO — Against Blue mold, use 2 to 3 lbs. per 100 gallons. Apply directly to plant beds at 3 to 6
gallons of spray per 100 sq. yard bed depending on size of plants. Start applications when leaves are about
the size of a dime, or when blue mold first appears. Apply at 3 to 4 day intervals as needed.

GLADIOLUS — Against Botrytis blight, Curvularia, and Stemphylium, use 1 1/2 pounds per 100 gallons of
water. Apply when diseases first appear, repeat at 7 day intervals. Shorten interval between applications to
3 to 5 days under severe disease conditions.

NOTE: The U. S. Food and Drug Administration has set tolerances for pesticide residues. Crops having residues in excess of
their permitted tolerances should not be marketed and to do so may violate State law and will violate Federal law when
sold in interstate commerce. Such crops should not be marketed without removal of such excess residues.

CAUTION

May cause irritation of nose, throat, and skin. Do not breathe dust or spray mist. May be harmful if inhaled or swallowed.
Avoid contact with skin, eyes, and clothing. Keep away from fire or sparks. In case of contact, flush with plenty of water.
for eyes, get medical attention. Keep containers closed. Store in dry place. Do not re-use container. Destroy when empty.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. The manufacturer makes no warranties,
expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all direc-
tions carefully.

DRAFT F 71 7283

Manufactured by **FLORIDA AGRICULTURAL SUPPLY COMPANY**, Jacksonville Florida, U.S.A.

FORM 700

ACCEPTED

9/1/55

NET CONTENTS
5 LBS.

ZINEB-75 WP

PRODUCT
No. 978

ACTIVE INGREDIENTS:

Zineb (Zinc ethylenebisdithiocarbamate)
(Zinc expressed as metallic 17.7%)

BY WT.
75.00%

INERT INGREDIENTS

25.00%

® Trade Marks Reg. U. S. Patent Office

TOTAL — 100.00%

CAUTION: KEEP OUT OF THE REACH OF CHILDREN.
See below for additional cautions.

USDA Reg. No. 2342-716

FASCO ZINEB-75 WP is a wettable powder fungicidal product containing 75% ZINEB fungicide. It is designed for use in aqueous sprays against certain diseases attacking vegetable, fruit, and ornamental crops as described below.

DIRECTIONS

Use contrary to these directions may cause serious plant, crop or personal injury, unsatisfactory control, or other unintended consequences.

CITRUS (Oranges, Grapefruit, Tangerines) — To control Rust Mite and associated russeting and greasy spot, use 1 pound per 100 gallons of water. Apply with very thorough coverage. Make application from late June through early August or earlier if rust mite begins to appear.

TOMATOES — Against Early and Late Blight, Septoria leaf spot, Gray Leaf spot (Stemphylium), Anthraco-
nose, use 1½ to 2 pounds per 100 gallons of water. Apply when disease is first present and repeat ap-
plications at 7 to 10 day intervals or oftener if blight conditions are severe. Do not exceed 4 lbs. per
acre.

PEPPERS, EGGPLANT — Against Alternaria, Anthraco-
nose, and Phomopsis of eggplant; Anthraco-
nose and
Cercospora leaf spot of pepper, use 1½ to 2 pounds per 100 gallons of water. Start applications at first
sign of disease and repeat at 7 to 10 day intervals as needed. Do not exceed 4 lbs. per acre on Eggplant;
8 lbs. per acre on Pepper.

CABBAGE, BROCCOLI, CAULIFLOWER — Against Downy Mildew and Alternaria, use 1½ to 2 pounds per
100 gallons of water. Apply at first sign of disease, repeat at 7 day intervals as needed. Do not apply
later than 7 days from harvest, do not exceed 6 lbs. per acre.

CANTALOUPE, CUCUMBERS, WATERMELONS, SQUASH — Against Downy Mildew, Anthraco-
nose, Gummy
Stem Blight, use 1½ to 2 pounds per 100 gallons of water. Apply at first sign of disease and repeat at 7
to 10 day intervals as long as necessary. Under severe disease conditions, shorten interval to 3 to 5 days.
Do not exceed 4 lbs. per acre.

TOBACCO — Against Blue mold, use 2 to 3 lbs. per 100 gallons. Apply directly to plant beds at 3 to 6
gallons of spray per 100 sq. yard bed depending on size of plants. Start applications when leaves are about
the size of a dime, or when blue mold first appears. Apply at 3 to 4 day intervals as needed.

GLADIOLUS — Against Botrytis blight, Curvularia, and Stemphylium, use 1½ pounds per 100 gallons of
water. Apply when diseases first appear, repeat at 7 day intervals. Shorten interval between applications to

plications at 7 to 10 day intervals or oftener if blight conditions are severe. Do not exceed 4 lbs. per acre.

PEPPERS, EGGPLANT — Against Alternaria, Anthracnose, and Phomopsis of eggplant. Anthracnose and Cercospora leaf spot of pepper, use 1½ to 2 pounds per 100 gallons of water. Start applications at first sign of disease and repeat at 7 to 10 day intervals as needed. Do not exceed 4 lbs. per acre on Eggplant; 8 lbs. per acre on Pepper.

PEPPER PEPPERS

CABBAGE, BROCCOLI, CAULIFLOWER — Against Downy Mildew and Alternaria, use 1½ to 2 pounds per 100 gallons of water. Apply at first sign of disease, repeat at 7 day intervals as needed. Do not apply later than 7 days from harvest, do not exceed 6 lbs. per acre.

CANTALOUPE, CUCUMBERS, WATERMELONS, SQUASH — Against Downy Mildew, Anthracnose, Gummy Stem Blight, use 1½ to 2 pounds per 100 gallons of water. Apply at first sign of disease and repeat at 7 to 10 day intervals as long as necessary. Under severe disease conditions, shorten interval to 3 to 5 days. Do not exceed 4 lbs. per acre.

TOBACCO — Against Blue mold, use 2 to 3 lbs. per 100 gallons. Apply directly to plant beds at 3 to 6 gallons of spray per 100 sq. yard bed depending on size of plants. Start applications when leaves are about the size of a dime, or when blue mold first appears. Apply at 3 to 4 day intervals as needed.

GLADIOLUS — Against Botrytis blight, Curvularia, and Stemphylium, use 1½ pounds per 100 gallons of water. Apply when diseases first appear, repeat at 7 day intervals. Shorten interval between applications to 3 to 5 days under severe disease conditions.

NOTE: The U. S. Food and Drug Administration has set tolerances for pesticide residues. Crops having residues in excess of their permitted tolerances should not be marketed and to do so may violate State law and will violate Federal law when sold in interstate commerce. Such crops should not be marketed without removal of such excess residues.

CAUTION

May cause irritation of nose, throat, and skin. Do not breathe dust or spray mist. May be harmful if inhaled or swallowed. Avoid contact with skin, eyes, and clothing. Keep away from fire or sparks. In case of contact, flush with plenty of water: for eyes, get medical attention. Keep containers closed. Store in dry place. Do not re-use container. Destroy when empty.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

DRAFT F

71-7283

Manufactured by **FLORIDA AGRICULTURAL SUPPLY COMPANY**, Jacksonville, Florida, U.S.A.

FORM 700

