



Net Contents  
50 Pounds

# COPPER SEVIN<sup>®</sup> SULPHUR DUST 5-5-75

Product No.  
3-1145-1

(FOR PEANUTS)

FASCO COPPER SEVIN<sup>®</sup> SULPHUR DUST 5-5-75 is a combination insecticide-fungicide dust containing SEVIN insecticide, Copper, and Sulphur for use on peanuts as described below.

### DIRECTIONS

PEANUTS — Against Spotted Cucumber Beetle (Southern Corn Rootworm), Thrips, Velvet Bean Caterpillar, Corn Earworm (Bollworm), Armyworm, Leafhoppers, and Leafspot apply 20 lbs. per acre when insects or disease first appear. Repeat at 7 to 10 day intervals as needed. Do not exceed 30 lbs. per acre. May be applied up to day of harvest.

If this product is used in accordance with label directions, the above crop may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk or meat.

NOTE: The U.S. Food and Drug Administration has set tolerance for pesticide residues. Crops having residues in excess of their permitted tolerances should not be marketed and to do so may violate State law and will violate Federal law when sold in interstate commerce. Such crops should not be marketed without removal of such pesticide residues.

ACTIVE INGREDIENTS	BY WT.
Carbaryl (1-naphthyl N-Methylcarbamate)	5.00%
Copper (Derived from Brown Copper Oxide)	5.00%
Sulphur	75.00%
INERT INGREDIENTS	15.00%
TOTAL	100.00%

Mesh Test: Note less than 90% through a 325 mesh screen.

U.S. Patent No. 2,963,478

Sevin<sup>®</sup> Trade Mark of UCC for 1-naphthyl N-methylcarbamate.

USDA Reg. No. 2342-727

**CAUTION: Keep Out of the Reach of Children. See additional cautions on this label.**

### CAUTION

HARMFUL IF SWALLOWED OR INHALED. Do not breathe dust. Do not take internally. Skin contact may be harmful. Avoid contact. Wash hands before eating or smoking. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of feedstuffs. Avoid contamination of feed, feeding troughs, and water receptacles. Sulphur may burn foliage when temperatures are extremely high. Do not make sulphur applications at such times. Do not apply when weather conditions favor drift of dust from areas treated.

### CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed.

The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read and direct use carefully.