


DANGER—POISON—PRECAUTIONS 


READ LABEL BEFORE USING I



POISONOUS IF SWALLOWED



POISONOUS BY SKIN CONTACT



POISONOUS IF BREATHED

POISONOUS TO FISH AND WILDLIFE

See Antidote and
Precautions on
this label



CAN CAUSE
DEATH

Keep Out of
Reach of
Children

Net Contents 50 Pounds

Product No. 3-1148-1

PARATHION DUST-1

ACTIVE INGREDIENTS

Parathion (0.0 Diethyl O p nitrophenyl Thiophosphate)

BY WT.

1.00%

INERT INGREDIENTS

99.00%

TOTAL - 100.00%

Handwritten notes:
DUST-1 - 1971
2nd - 750

CROP USE DIRECTIONS

FASCO PARATHION DUST 1 is an insecticidal dust for use against certain insects attacking economic crops as directed below

DIRECTIONS

Use contrary to these directions may cause serious crop plant or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

BEANS - Against Mexican Bean Beetle, Aphids, Leafhoppers, Lygus and Stink bugs, dust thoroughly with 25 to 40 pounds (no more) per acre every 7 to 10 days or as needed, beginning when insects first appear. Do not apply within 15 days of harvest.

CABBAGE CAULIFLOWER BROCCOLI COLLARDS MUSTARD - Against Aphids, Diamond Back Moth Larvae, Imported Cabbageworm and Armyworm, use at the rate of 25-40 lbs. per acre every 3 to 10 days if needed. Do not use on Cauliflower and Broccoli within 7 days, Cabbage within 10 days, Collards and Mustard within 14 days before harvest.

CUCUMBER SUMMER SQUASH - Against Striped and Spotted Cucumber Beetle, Aphids, Flea Beetle, Leaf Miners, Pickworms and Melonworm, dust thoroughly with 25-40 lbs. (no more) per acre every 7 to 10 days, beginning when insects first appear. Against Squash Bugs, apply 50 lbs. per acre when insects first appear. Repeat as needed. Do not treat cucumbers before vining.

Do not apply within 15 days of harvest
TOMATO EGGPLANT PEPPER - Against Flea Beetles, Aphids, Spider Mites, Leaf Miners, dust thoroughly with 25 to 40 pounds per acre (no more) every 7 to 10 days. Do not apply to tomatoes within 10 days of harvest, nor to Eggplant or Pepper within 15 days of harvest.

POTATO - Against Colorado Potato Beetle, Aphids, Leaf Miner, Flea Beetles, dust thoroughly with 25 to 40 pounds per acre at 7 to 10 day intervals. Do not apply within 5 days of harvest.

TOBACCO - Against Aphids, Stinkbugs, and Flea Beetles, apply 15-30 pounds per acre every 7 to 10 days, beginning when insects first appear.

CAUTION - Do not apply to tobacco within 5 days of pruning or 15 days of cutting. Handlers should wear gloves where applications have been made within 30 days of cutting. Avoid plant juices coming in contact with skin or other parts of the body when engaged in cutting the crop.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

NOTE TO PHYSICIAN

Parathion is a highly effective insecticide. It is also a powerful poison. It is absorbed through the skin, lungs, and gastrointestinal tract. Symptoms of poisoning include nausea, vomiting, diarrhea, muscle twitching, and respiratory distress. In severe cases, it can lead to respiratory failure and death. Treatment should be initiated immediately upon suspicion of poisoning.

POST-TREATED AREA

Do not enter treated areas for 24 hours after application. Avoid contact with treated surfaces.

CONTAINER DISPOSAL

Do not reuse or refill this container. Dispose of empty containers in accordance with local regulations. Do not burn or incinerate.

FIRST AID TREATMENT

If breathing has stopped, give artificial respiration. If swallowed, do not induce vomiting. If on skin, wash with soap and water. If in eyes, flush with water for 15 minutes. Seek medical attention immediately.