



Net Contents
50 Pounds

COPPER SEVIN[®] SULPHUR DUST 4-5-75

Product No.
3-1157-1

FASCO COPPER SEVIN[®] SULPHUR DUST 4-5-75 is a combination insecticide-fungicide dust containing Copper, Sevin insecticide and Sulphur for use on peanuts as described below.

DIRECTIONS

PEANUTS — Against Spotted Cucumber Beetle (Southern Corn Rootworm), Thrips, Velvet Bean Caterpillar, Corn Earworm (Bollworm), Armyworms, Leafhoppers, and Leafspot apply 25 lbs. per acre when insects or disease first appear. Repeat at 7 to 10 day intervals as needed. Do not exceed 30 lbs. per acre. May be applied up to day of harvest.

If this product is used in accordance with label directions, the above crop may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk or meat.

NOTE: The U.S. Food and Drug Administration has set tolerances for pesticide residues. Crops having residues in excess of their permitted tolerances should not be marketed and to do so may violate State law and will violate Federal law when sold in interstate commerce. Such crops should not be marketed without removal of such excess residues.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy it by burying with waste or burning. Stay out of smoke and fumes.

(FOR PEANUTS)

ACTIVE INGREDIENTS

Copper (Derived from Brown Copper Oxide)	4.00%
Carbaryl (1-naphthyl N-Methyl carbamate)	5.00%
Sulphur	75.00%

INERT INGREDIENTS

16.00%

Total — 100.00%

Sevin[®] Trade Mark of UCC for 1-naphthyl N-methyl carbamate.
Inert Test: Not less than 90% through a 325 mesh screen.

U.S. Patent No. 2,933,471

USDA Reg. No. 1342-731

CAUTION: Keep Out of the Reach of Children. Note other Cautions on this label.

CAUTION

HARMFUL IF SWALLOWED OR INHALED. Do not breathe dust. Do not take internally. Skin contact may be harmful. Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs, and watering receptacles. Sulphur may burn foliage when temperature is high. Do not make sulphur applications at such times. Do not apply when weather conditions favor drift of dust from crop areas treated. Birds feeding on treated areas may be killed.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed.

The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.