



12/09/71

2342-745

NET CONTENTS
50 LBS.

THIODAN® ZINEB DUST 3-6

PRODUCT
No. 777991

FASCO THIODAN-ZINEB DUST 3-6 is an insecticidal fungicidal dust formulation which contains 3% THIODAN insecticide and 6% Zineb fungicide. This product is designed for control of certain insects and diseases attacking vegetable crops as directed below.

DIRECTIONS

Use contrary to these directions may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

CUCUMBER, SUMMER SQUASH, WINTER SQUASH, MELONS — Against Anthracnose, Downy Mildew, Gummy stem blight, Aphids, Cucumber Beetles, and Squash vine borer, apply 25-35 pounds per acre commencing when first signs of disease appear or when insects first appear. Apply at 7 day intervals. May be applied up to day of harvest.

MELONS — Against Cabbage Looper, Omnivorous leaf roller, Aphids, Cucumber beetle, Anthracnose, Downy Mildew, and Gummy stem blight, apply 35-50 lbs. per acre when insects or disease first appear. Repeat at weekly intervals. Do not exceed 50 lbs. per acre. May be applied up to day of harvest.

TOMATO, PEPPER — Against Early and Late Blight, Frogeye Spot of Pepper, Aphids, Flea beetle, Colorado potato beetle, Tomato fruitworm, and Hornworm, apply 33 pounds per acre. Begin applications when insects or disease first appear, repeat every 3 to 7 days as needed. Do not exceed 33

DRAFT D 71-8194

ACTIVE INGREDIENTS:	By Wt.
Endosulfan*	3.00%
Zineb (Zinc ethylene bisdithiocarbamate) (Equivalent to Zinc, expressed as metallic 1.42%)	6.00%
INERT INGREDIENTS	91.00%
TOTAL	100.00%

*Hexachlorocyclohexane-2,4,3-benzodioxathiepin oxide
Thiodan is a registered trademark of Farbwerke Hoechst A.G.

U.S. Patent No. 2,799,685
EPA Reg. No. 2342-745

CAUTION: Keep Out of the Reach of Children. See additional cautions on this label.

pounds per acre. Do not apply within 1 day of harvest.

EGGPLANT — Against Phomopsis Blight, Alternaria leaf spot, Aphids, Colorado potato beetle, and Flea beetle, apply 20 lbs. per acre when disease or insects first appear. Repeat at 7 day interval or as needed. Do not exceed 20 lbs. per acre. Do not apply within 1 day of picking.

BROCCOLI, CABBAGE — Against Cabbage Aphids, Cabbage looper, Imported cabbageworm, Diamond-back moth larvae, and Alternaria leaf spot, apply 25-33 lbs. per acre. Begin applications when insects or disease first appear, repeat every 5 to 7

days as needed. Do not exceed 33 lbs. per acre. Do not apply within 7 days of harvest.

NOTE—Crops having residues in excess of their permitted tolerances should not be marketed and to do so may violate State law and will violate Federal law when sold in interstate commerce. Such crops should not be marketed without removal of such excess residues.

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of feed and foodstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Workers entering treated areas on the day of application should wear protective clothing.

DANGER TO FISH AND WILDLIFE

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Treated seeds exposed on soil surface may be hazardous to birds and other wildlife. Apply this product only as specified on this label.

CONTAINER DISPOSAL

Do not re-use empty container. Destroy when empty by burying with wastes well away from water supplies or by burning. (Stay well away from smoke and fumes).

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

Manufactured by **KERR-MCGEE CHEMICAL CORP.** Jacksonville, Florida, U.S.A. General Offices Oklahoma City, Oklahoma