

COPPER SPRAY

Gen. Use: Copper Spray is a neutral copper spray concentrate for use against a wide range of fungal and bacterial diseases of plants.

DIRECTIONS: Use contrary to these directions may cause injury to plants, crop or persons, unsatisfactory control or other unintended consequences.

Citrus: Against Scab and/or Melanose, spray twice, once in late winter (Dormant spray) just before new growth begins and repeat when $\frac{2}{3}$ of the petals have fallen (post bloom spray).

For Melanose alone, spray 1 to 3 weeks after petals fall. A second application may be made 4 weeks later if the disease has been troublesome, and if the bloom is late or scattered such as during a wet spring. For post bloom spray, mix 1 tablespoon per gal. of water, $\frac{1}{4}$ cup for 5 gal. or $\frac{1}{2}$ cup per 10 gal. For dormant spray, double rate.

Flowers, Shrubs: Against leaf spot and other diseases such as Black Spot of Rose, Abrusvite B'ght, False Anthracnose, spray to wet all foliage and above ground parts at the first sign of disease using 3 table-
spoons per gal. of water, $\frac{1}{2}$ cup per 5 gal. or 1 cup per 10 gal. Repeat as needed.

Grasses: Against Gray Leafspot (Panicum) and St. Augustine Grass, use $1\frac{1}{2}$ cups per 10 gal. of water per 15 to 20 gal. of spray and apply to 1000 sq. ft.

Repeat at the first sign of disease and repeat every 10 to 14 days as needed.

Tomatoes, Peppers, Potatoes: Against Early Spot Disease of Tomato and Pepper, Frogeye Spot of Pepper and Early and Late Blights of Tomato and Potato, spray at the first sign of disease using Flower and Shrub rate above. Repeat every 5 to 10 days as needed.

CAUTION: Harmful if swallowed! If taken internally, induce vomiting by giving mustard and water and tickling the throat. Give large doses of magnesium or lime water, followed by doses of Epsom Salts.

Danger To Fish and Wildlife: This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Container Disposal: Do not reuse empty container. Destroy it by puncturing or crushing. Bury or discard in a safe place away from water supplies.

CONDITIONS OF SALE: All statements concerning the use of this product apply only when used as directed. The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

VARAD-TINE

POSITIVE PESTICIDES

COPPER SPRAY

For disease of
citrus, flowers,
shrubs, vegetable


ACTIVE INGREDIENT	BY WT.
Copper complexed with Malic	95.0%
(Derived from the Copper Sulphate)	1.0%
INERT INGREDIENTS	4.0%

CAUTION: Keep out of reach of children. See back panel for additional cautions.

Net Contents 1 1/2 Pounds

Manufactured by Kerr-McGee Chemical Corp.
Jacksonville, Florida U. S. A.
General Offices, Oklahoma City, Oklahoma

To control LICHENS on trunks and branches of shrubs and trees, spray to thoroughly wet LICHENS, using 1 tablespoon per gallon of water. Repeat as needed.



REGISTRATION OFFICE
PESTICIDES
12 APR 1961