

DIRECTIONS: For control of Brown Patch (Rhizoctonia Solani) attacking St. Augustine, Bermuda and similar grasses, apply 2 tablespoons per gal. of water to 30 sq. ft. (5 ft. x 6 ft.) or 1 pint in 15 to 20 gal, water to cover 500 sq. ft. (20' x 25'). For easy application, use hose-end type sprayer designed to deliver 15 to 20 gal, dilute spray while emptying qt. jar. Pour 1 pint concentrated liquid into jar and fill to top with water and apply to 500 sq. ft, Apply in spring or fall at first sign of disease and repeat in 3 to 4 weeks if necessary. Sprinkle after application to wash spray into soil surface. Follow recommended rate. Excess may cause a temporary yellowing FISH AND WILDLIFE CAUTION: This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equip-

ment, or disposal of wastes.

CAUTION: Avoid inhaling spray mist. This material may cause skin irritation. Wash hands and face thoroughly after using. Do not use or store near heat or open flame. Use adequate ventilation. Keep container closed. Avoid contamination of feed and foodstuffs. Do not graze treated areas. Do not feed clippings to livestock. Keep out of reach of children. Do not re-use empty container. Wash thoroughly with water and detergent. Discard in a safe place. **CONDITIONS OF SALE:** All statements concerning the use of this product apply only when used as directed. The manufacturer makes no warranties, expressed

or implied, concerning this product or

its use, which extend beyond the de-

scription on the label. Read all directions

carefully.



