




DANGER—POISON—PRECAUTIONS 


POISONOUS IF SWALLOWED

This product can kill you if swallowed even in small amounts; spray mist or dust may be fatal if swallowed.



POISONOUS BY SKIN CONTACT

Poisonous if touched by hands or spilled or splashed on skin, in eyes or on clothing; liquid goes through clothes.



POISONOUS IF BREATHED

Breathing vapors, spray mist or dust may be fatal.

POISONOUS TO FISH AND WILDLIFE

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Manufactured by
KERR-MCGEE CHEMICAL CORP.
Jacksonville, Florida, U.S.A.
General Offices Oklahoma City Oklahoma

STOP READ LABEL BEFORE USING!

POISON

See Side Panels
for Antidote &
Precautions



DANGER

Keep Out of
Reach of
Children

TOX-MP 4-4 EC

COTTON AND SOYBEAN INSECTICIDE SPRAY

ACCEPTED
APR 30 1979

ACTIVE INGREDIENTS:

- O,O dimethyl O,p nitrophenyl thiophosphate
- Toxaphene (Chlorinated Camphene)
(Chlorine Content 67% to 69%)
- Xylene Range Solvent

INERT INGREDIENTS

- Contains 4 lbs. Actual Toxaphene plus 4 lbs. Actual Methyl Parathion per Gallon
- Methyl Parathion

BY WT.

- 37.30%
- 37.30%
- 18.75%
- 6.65%

TOTAL 100.00%

NOT FOR HOME USE

NET CONTENTS
GALLONS

FASCO

PRODUCT NO
20-77970

FASCO TOX MP 4-4EC is a water emulsifiable insecticidal concentrate designed for use against certain insect pests attacking economic crops described on this label.

USE IN POWER OPERATED GROUND SPRAYERS
Use specified dosage of the spray concentrate in 5-10 gallons of water (low volume) to 50-100 gallons (high volume) per acre.

USE IN AERIAL APPLICATION Use specified dosage of the spray concentrate in 3-10 gallons water per acre.

DIRECTIONS

Use contrary to these directions may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences. For best results premix with a small amount of water and mix well before addition to spray tank.

COTTON (Early Season) Apply when needed at rate of 1 Quart per acre (1 Gallon per 4 acres) against overwintered Bollweevils, Aphids, Thrips, and Cotton Fleahoppers. (Mid and Late Season) Apply when needed at rate of 1 1/2 Quarts per acre against Boll Weevils, Aphids, Cotton Bollworm, Spider Mites, Thrips, Cotton Fleahoppers, Cotton Leafworms, and Lygus Bugs. Do NOT apply within 7 days of hand picking or harvest. Workers entering the fields within 24 hours after application should wear protective clothing. Do not graze dairy animals or animals being finished for slaughter on fields treated late in the season.

SOYBEANS Against Armyworms, Velvetbean caterpillars, Corn earworm, Blister beetles, Mexican Bean Beetle, and Stink bugs, apply 1 Quart per acre (1 gallon per 4 acres) in sufficient water for good coverage. Make application when insects first appear, repeat as needed. Do not apply more than two applications per growing season. Do not apply within 21 days of harvest. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals, or animals being finished for slaughter. Do not feed soybean meal trash to livestock or poultry. Workers entering the field within 24 hours after application should wear protective clothing.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

ACQU...
4/30/73

EPA Reg. No. 2342-773

2342-773

4/30/73

USE ONLY WHEN WEARING THE FOLLOWING
PROTECTIVE EQUIPMENT AND CLOTHING

Wear eye protection and clothing that will protect you from splashes. Do not use in confined spaces. Do not use near open flames or heat. Do not use if you are pregnant or nursing. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease or other cardiovascular conditions. Do not use if you have a history of kidney disease or other renal conditions. Do not use if you have a history of liver disease or other hepatic conditions. Do not use if you have a history of diabetes or other endocrine conditions. Do not use if you have a history of epilepsy or other neurological conditions. Do not use if you have a history of mental health conditions. Do not use if you are taking any medications. Do not use if you are drinking alcohol. Do not use if you are under the influence of any drugs. Do not use if you are feeling dizzy, lightheaded, or otherwise unwell. Do not use if you are pregnant or nursing. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease or other cardiovascular conditions. Do not use if you have a history of kidney disease or other renal conditions. Do not use if you have a history of liver disease or other hepatic conditions. Do not use if you have a history of diabetes or other endocrine conditions. Do not use if you have a history of epilepsy or other neurological conditions. Do not use if you have a history of mental health conditions. Do not use if you are taking any medications. Do not use if you are drinking alcohol. Do not use if you are under the influence of any drugs. Do not use if you are feeling dizzy, lightheaded, or otherwise unwell.

WORK SAFETY RULES

Do not use in confined spaces. Do not use near open flames or heat. Do not use if you are pregnant or nursing. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease or other cardiovascular conditions. Do not use if you have a history of kidney disease or other renal conditions. Do not use if you have a history of liver disease or other hepatic conditions. Do not use if you have a history of diabetes or other endocrine conditions. Do not use if you have a history of epilepsy or other neurological conditions. Do not use if you have a history of mental health conditions. Do not use if you are taking any medications. Do not use if you are drinking alcohol. Do not use if you are under the influence of any drugs. Do not use if you are feeling dizzy, lightheaded, or otherwise unwell.

PREVENTION

Do not use in confined spaces. Do not use near open flames or heat. Do not use if you are pregnant or nursing. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease or other cardiovascular conditions. Do not use if you have a history of kidney disease or other renal conditions. Do not use if you have a history of liver disease or other hepatic conditions. Do not use if you have a history of diabetes or other endocrine conditions. Do not use if you have a history of epilepsy or other neurological conditions. Do not use if you have a history of mental health conditions. Do not use if you are taking any medications. Do not use if you are drinking alcohol. Do not use if you are under the influence of any drugs. Do not use if you are feeling dizzy, lightheaded, or otherwise unwell.

FIRST AID TREATMENT

Do not use in confined spaces. Do not use near open flames or heat. Do not use if you are pregnant or nursing. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease or other cardiovascular conditions. Do not use if you have a history of kidney disease or other renal conditions. Do not use if you have a history of liver disease or other hepatic conditions. Do not use if you have a history of diabetes or other endocrine conditions. Do not use if you have a history of epilepsy or other neurological conditions. Do not use if you have a history of mental health conditions. Do not use if you are taking any medications. Do not use if you are drinking alcohol. Do not use if you are under the influence of any drugs. Do not use if you are feeling dizzy, lightheaded, or otherwise unwell.

HAZARD IDENTIFICATION

Do not use in confined spaces. Do not use near open flames or heat. Do not use if you are pregnant or nursing. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease or other cardiovascular conditions. Do not use if you have a history of kidney disease or other renal conditions. Do not use if you have a history of liver disease or other hepatic conditions. Do not use if you have a history of diabetes or other endocrine conditions. Do not use if you have a history of epilepsy or other neurological conditions. Do not use if you have a history of mental health conditions. Do not use if you are taking any medications. Do not use if you are drinking alcohol. Do not use if you are under the influence of any drugs. Do not use if you are feeling dizzy, lightheaded, or otherwise unwell.

ENVIRONMENTAL PRECAUTIONS

Do not use in confined spaces. Do not use near open flames or heat. Do not use if you are pregnant or nursing. Do not use if you have a history of asthma or other respiratory conditions. Do not use if you have a history of heart disease or other cardiovascular conditions. Do not use if you have a history of kidney disease or other renal conditions. Do not use if you have a history of liver disease or other hepatic conditions. Do not use if you have a history of diabetes or other endocrine conditions. Do not use if you have a history of epilepsy or other neurological conditions. Do not use if you have a history of mental health conditions. Do not use if you are taking any medications. Do not use if you are drinking alcohol. Do not use if you are under the influence of any drugs. Do not use if you are feeling dizzy, lightheaded, or otherwise unwell.

DANGER TO LIFE

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME