

DIBEN[®] SPRAY

INSECTICIDE

DIRECTIONS FOR USE

Apply as a thorough cover spray. Make applications when insects first appear or in advance of an expected infestation. Repeat applications as needed or as noted below.

Use coarse drenching sprays for lawns, soil, and other surface applications. Use clean sprayer and carefully measure and mix correct amount with water. Use medium-fine mist spray for trees and plants. Flush sprayer with clean water after each use. Do not use on certain ferns, poinsettia, hibiscus, papaya, pilea, gardenia and orchids since injury may occur.

LAWNS (Bluegrass, Merion, Fescue, St. Augustine, Dichondra, Bent, Bermuda): To control Chinch Bugs, Sod Webworms, Bermuda Mites, Ants, Ticks, Earwigs & Lawn Billbugs use 2 tablespoons (1 fluid ounce) in 3 gallons of water per 125 square feet, 8 fluid ounces in 24 gallons of water per 1,000 square feet. Thoroughly wet down grass a few hours before applying. For Lawn Billbugs water lawn thoroughly after treatment.

ROSES AND FLOWERS (Roses, Azaleas, Chrysanthemums, Gladioli, Carnations): To control Aphids, Mites, Leafhoppers, and Thrips use 2 tablespoons (1 fluid ounce) per 3 gallons of water. Apply as a thorough foliar spray. Spray underside of leaves. Wet plants to the dripping point.

TREES AND SHRUBS (Oak, Maple, Locust, Holly, Boxwood, Pine, Spruce, & Willow): To control Aphids, Mites, Scale Crawlers, Leaf Miners & Bagworms use 2 tablespoons (1 fluid ounce) per 3 gallons of water. Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to penetrate dense foliage.

FRUITS (Apples, Peaches, Oranges, Plums, Grapes, Cherries & Strawberries): To control Codling Moths, Mites, Aphids, Leaf Rollers & Scale Crawlers use 2 tablespoons (1 fluid ounce) per 3 gallons of water. Apply as a full cover spray. Start spraying after blooming and repeat at 10 to 14 day intervals if needed. Do not spray fruit within 21 days of picking.

VEGETABLES (Tomatoes, Carrots, Potatoes, Beans, Cucumbers, Onions, Cantaloupes, Peppers & Cabbage) use 2 tablespoons (1 fluid ounce) in 3 gallons of water per 1,000 square feet. Generally application at 7-10 day intervals may be required. Do not apply within one day of picking tomatoes, 35 days of harvesting potatoes, or 14 days of picking all other crops listed in this section.

*To control **Bean Beetles, Aphids, Leaf Miners, Imported Cabbageworms, Cucumber Beetles**

NUISANCE PESTS IN OUTSIDE AREAS: To control Elias, Ants, Cockroaches, Boxelder Bugs, Clover Mites, Spiders, Ticks, Chiggers, & Earwigs, use 8 fluid ounces per 3 gallons of water. Thoroughly spray window frames, door sills, screens, porch and patio walls and ceilings, foundations, walks, garbage cans, and into cracks and other places where insects may hide. Do not treat plants or animals with this dosage rate.

IMPORTANT: Do not use on household pets or humans. Do not permit children or pets to go onto the sprayed grass until the spray has completely dried. Do not use in the home.

PRECAUTIONARY INFORMATION: WARNING! May be fatal if swallowed. May be harmful if inhaled or absorbed through skin.

Do not breathe vapor or spray mist. Avoid contact with eyes, skin and clothing. Keep container closed. Use with adequate ventilation. Avoid storage near feed or food products. Wash thoroughly after handling. Wash clothing before reuse. Do not use, pour, spill or store near heat or open flame.

FIRST AID: If swallowed, induce vomiting by sticking finger down throat or by giving soapy or strong salty water to drink. Repeat until vomit is clear. Call a physician. Never give anything by mouth to an unconscious person. If inhaled, remove to fresh air. Call a physician. In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Call a physician.

DANGER TO FISH AND WILDLIFE: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL: Do not re-use the container. Destroy when empty.

CONDITIONS OF SALE: All statements concerning the use of this product apply only when used as directed. The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

GRO-TONE[®]

DIBEN[®] SPRAY

INSECTICIDE
CONTAINS DIAZINON†

CONTROLS

Ants, Aphids, Bagworms, Bean Beetles, Bermuda Mites, Boxelder Bugs, Chiggers, Chinch Bugs, Clover Mites, Cockroaches, Codling Moths, Cucumber Beetles, Earwigs, Flies, Imported Cabbageworms, Lawn Billbugs, Leafhoppers, Leaf Miners, Leaf Rollers, Scale Crawlers, Sod Webworms, Spiders, Ticks, Thrips.

APPLY TO

Certain Lawns, Vegetables, Fruits, Outside Areas.
See Back Panel For Specific Use Directions.

ACTIVE INGREDIENTS	BY WEIGHT
0, 0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	25%†
Aromatic Petroleum Derivative Solvent	56%
INERT INGREDIENTS	19%

† DIAZINON INSECTICIDE, DIAZINON, REGISTERED TRADEMARK, CIBA-GEIGY CORPORATION, U.S. PATENT NO. 2,754,243.

WARNING: KEEP OUT OF REACH OF CHILDREN
See back panel for additional precautionary information.

Net Contents 1 Quart

EPA REG NO. 2342-777
C-12-73-W

PRODUCT NO 725812

REVISION B

MANUFACTURED BY
 **KERR-MCGEE CHEMICAL CORP.**
OKLAHOMA CITY, OKLAHOMA

