



4-80 PEANUT DUST

U.S.D.A. Reg. No. 9132-4

ACTIVE INGREDIENTS:

Copper expressed as metallic* 1.00%

Sulfur 80.00%

INERT INGREDIENTS:

16.00%

100.00%

*Copper derived from cuprous and cupric oxides.

See additional information elsewhere on label

See additional information elsewhere on label


READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

Burn bag immediately when empty. Stay out of smoke.

Harmful if swallowed. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur is safe in that locality. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

DIRECTIONS

PEANUTS: Early and Late Leaf Spot Infections— Apply 20 to 25 pounds of 4-80 Peanut Dust per acre at 7 to 10-day intervals or as necessary. Begin dusting when diseases first appear and continue to within 1 month of harvest.



KERR-MCGEE CHEMICAL CORP.



GENERAL OFFICES: OKLAHOMA CITY, OKLAHOMA
NEW BERN, NORTH CAROLINA • CHARLESTON, SOUTH CAROLINA