

METHYL TRITHION 4 - E

EMULSIFIABLE LIQUID AN ORGANIC PHOSPHATE INSECTICIDE-ACARICIDE (Contains 46.5% Technical Methyl Trithion)






Keep out of reach of children.

May be fatal if swallowed, inhaled, or absorbed through skin. Do not get in eyes or on skin or on clothing. If material gets in eyes, flush with water for 15 minutes. In case of skin contact, wash immediately with soap and water. Do not breathe spray mist. For prolonged commercial application, wear rubber gloves, protective clothing and rubber overshoes. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothes with soap and hot water before re-use. Do not contaminate food and feed stuffs. Use this product at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. Do not apply on cotton after half the bolls are open. Do not graze or feed treated trash to livestock.

This product is poisonous to tish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label.

Methyl Trithion 4-E is an emulsifiable liquid for use in water as an insecticidal spray. Apply when air is calm, using ground equipment or aircraft. Make thorough coverage of plants. Apply at the first sign of infestation and repeat at regular intervals when needed to maintain control.

Pour specified amount of this product into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. If mixture does not mix readily, but tends to separate as an oily layer, do not use, as injury to plants may result. Do not combine with wettable powders unless previous use of the mixture has been proven physically compatible and safe to plants. Always thoroughly emulsify this product with at least half of total water before adding wettable powder.





ACTIVE INGREDIENTS:

*S—[[(p-chloropheny!)thio[mothyl] 0, 0-dimethyl phosphorodithioate *Related Organic Phosphates Xylene Range Aromatic Solvent

INERT INGREDIENTS:

RECOMME

37.29

7.09

42.09

13.8%

100.0%

DIRECTIONS FOR USE

The rates given are to be diluted w type of equipment to be used. The such a manner to provide good cov

Cotton: Early Season-Cotton aphids bugs, over-wintering boll weevil, sp acre in sufficient water to provide Season-Boll weevil, cotton aphid, bugs, spider mites, thrips. Apply 1 provide adequate coverage. Repea heavy migration of boll weevil or h every 3 to 5 days.

*From 46.5% Technical Methy! Tri

Keep away from heat and open tures 32° F*

*Liquid phases separate at lower higher temperature and mix thorou assure uniformity before use.

ا الا الا الا الحالية التي المركزة المحلية المركزة المحلية المحلية المحلية المحلية المحلية المحلة المحلة المحل المحلية المحلومة المحلية المحلية المحلية المحلية المحلية المحلية المحلية المحلية المحلية المحلة المحلية المحلية ومحلي محلة المحلومة المحلومة المحلومة المحلية المحلية المحلية المحلية المحلية المحلية المحلية المحلية المحلية ا ERR-MCGEE CHEMICAL CORP.

THYL TRITHION 4 - E

AN ORGANIC PHOSPHATE FIABLE LIQUID INSECTICIDE-ACARICIDE

(Contains 46.5% Technical Methyl Trithion)

Keep out of reach of children.

absorbed through skin. Do ig. If material gets in eyes, se of skin contact, wash imbreathe spray mist. For prober gloves, protective clothing ms and face thoroughly with ing. Wash all contaminated re-use. Do not contaminate t at recommended rates and rvest than specified. Do not ed by unprotected humans or i, fiber or pasture crops. Do re open. Do not graze or feed

llife. Keep out of lakes, ponds not specified on the label.

DIRECTIONS FOR USE

Methyl Trithion 4-E is an emulsifiable liquid for use in water as an insecticidal spray. Apply when air is calm, using ground equipment or aircraft. Make thorough coverage of plants. Apply at the first sign of infestation and repeat at regular intervals when needed to maintain control


Pour specified amount of this product into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. If mixture does not mix readily, but tends to separate as an oily layer, do not use, as injury to plants may result. Do not combine with wettable powders unless previous use of the mixture has been proven physically compatible and safe to plants. Always thoroughly emulsify this product with at least half of total water before adding wettable powder.

Keep away from heat and open flame. Minimum storage temperatures 32° F*

*Liquid phases separate at lower temperatures. Warm or store at higher temperature and mix thoroughly to recombine liquid phases and assure uniformity before use.

at realities remaining an teat crops of the spont stream weak of the second end of t





ACTIVE INGREDIENTS:

*S-[[(p-chlorophenyl)thio]methyl] O.O-dimethyl phosphorodithioate *Related Organic Phosphates Xylene Range Aromatic Solvent

INERT INGREDIENTS:

O-anneuryi	37.2% 7.0% 42.0%	
	<u>13.8%</u> 100.0%	USDA Reg. No. 9132-19
		وقيم محمد بعد المراجع والمحمد المناقبة فالمناف المتحد والمتكفي الجرام محمو والمراج ومعطوا المدور ومعروا وال

RECOMMENDATIONS

The rates given are to be diluted with sufficient water, depending upon type of equipment to be used. The finished spray should be applied in such a manner to provide good coverage.

Cotton: Early Season-Cotton aphids, cotton leafworm, fleahopper, lygus bugs, over-wintering boll weevil, spider mites, thrips. Use ³/₄ pint per acre in sufficient water to provide adequate coverage. Mid and Late Season-Boll weevil, cotton aphid, cotton leafworm, fleahopper, lygus bugs, spider mites, thrips. Apply 1 pint per acre in sufficient water to provide adequate coverage. Repeat application every 5 to 7 days. If heavy migration of boll weevil or high populations are apparent, apply every 3 to 5 days.

*From 46.5% Technical Methyl Trithion