

FASCO

Net Contents
3 Pounds

ACCEPTED
THIRAM 75 WP

Product No.
3-1029-1

ACTIVE INGREDIENTS Thiram (Tetramethylthiuram disulfide)
INERT INGREDIENTS

BY WT.
75.00%
25.00%

TOTAL — 100.00%

CAUTION: Keep Out of the Reach of Children. See below for additional cautions.

USDA Reg. No. 2342-613

FASCO THIRAM 75 WP is a wettable powder fungicide formulation designed for the prevention and control of certain diseases attacking celery, tomato, and turf grass as described below.

DIRECTIONS

Use contrary to these directions may cause serious plant, crop or personal injury, unsatisfactory control or other unintended consequences.

CELERY — For control of early blight (Cercospora), late blight (Septoria) and Rhizoctonia, use 1 1/2 lbs. per 100 gallons water.

Plant Beds — Apply 15 to 25 gallons per 1200 Sq. Ft. of bed at 3-day intervals when disease first appears; shorten intervals under severe disease conditions.

Field — (Florida Only) Apply as full coverage spray (up to 200 gallons per acre) when disease appears. Repeat at weekly intervals or every 3-5 days under severe conditions. Do not apply within 7 days before harvest. Remove residues by stripping, trimming, and washing.

TOMATOES — (Florida Only) For the control of Gray Mold (Botrytis) and Rhizoctonia fruit rot, use 1 1/2 lbs. per 100 gallons of water. If gray leaf spot, late blight, or other foliage diseases appear, use in combination with the recommended fungicides. Make applications at 3 to 5 day intervals as long as disease threatens. Begin application after transplant or emergence and continue at 3-5 day intervals. Do not exceed 200 gallons of spray per acre.

TURF GRASSES (LAWNS, GOLF COURSES, PARKS, TURF NURSERIES) Against Brown patch (Rhizoctonia solani), Dollar Spot, Gray leaf spot, and Helminthosporium leaf spot, apply when disease first appears at the following rate:

Rate per 1,000 Square Feet 4 1/2 oz. (1 cupful) in 5 to 10 gallons of water
Rate per 5,000 Square Feet 1 1/2 lbs. in 25 to 50 gallons of water

Repeat applications at 7 to 10 day intervals as needed. Reduce intervals between sprays to 3 to 5 days when frequent rains may make conditions unusually favorable for disease development.

NOTE: Do not water or mow grass after treatment until the fungicide is thoroughly dry. Do not permit animals to feed or graze on treated area. Do not use grass clippings from treated areas for food or feed purposes. Do not re-use empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

DANGER TO FISH AND WILDLIFE

This product is toxic to fish. Keep out of lakes, streams and ponds. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes.

MAY CAUSE IRRITATION OF EYES, NOSE, THROAT, SKIN. HARMFUL IF INHALED OR SWALLOWED. Do not inhale dust or spray/mist. Do not get in eyes or on skin. Wash thoroughly after handling. In case of contact, flush with plenty of water; for eyes, get medical attention. Consumption of alcoholic beverages increases the possibility of harm.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

DRAFT C

71-6847

Manufactured by **KERR-MCGEE CHEMICAL CORP.** Jacksonville, Florida S.A.

800-L