

[2342-796]



4# METHYL PARATHION

Emulsifiable Concentrate

Contains 4 lbs. of Methyl Parathion per gallon

ACTIVE INGREDIENTS:

O,O-Dimethyl O-p-nitrophenyl thiophosphate
Xylene Aromatic Petroleum Solvent

INERT INGREDIENTS:

ACCOUNT
Jun-12-19

**POISON
DANGER**



**Keep out of reach
of children.**

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN. REPEATED INHALATION OR SKIN CONTACT MAY WITHOUT SYMPTOMS, PROGRESSIVELY INCREASE SUSCEPTIBILITY TO METHYL PARATHION POISONING. Do not swallow or get in eyes, on skin or on clothing. Do not breathe dust or vapors. Wear clean natural rubber gloves, clean waterproof protective clothing and goggles. Replace gloves frequently and destroy used gloves. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for methyl parathion protection. Wash thoroughly with soap and water after handling and before eating or smoking. If the material gets into the eyes immediately flush with water for at least ten minutes and get medical attention. In case of accidental contact, remove all contaminated clothing, and wash skin with soap and water; wash clothing before re-use. In case of spillage, wash down with large amounts of water. Workers entering fields within 24 hours of treatment should wear protective clothing.

This product is poisonous to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Unprotected human and animals should be kept out of treated fields on day of treatment. Keep out of reach of children and domestic animals. Decontaminate containers by washing thoroughly with a concentrated solution of alkali and detergent and rinse with water. Then destroy used containers to prevent reuse. Do not store near food or feed products. Keep away from heat and open flame.

ANTIDOTE AND FIRST AID TREATMENT: Atropine is the emergency antidote for Methyl Parathion poisoning. Consult your physician about obtaining an adequate supply of 1/100 grain atropine tablets for emergency use. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

INTERNAL: If the material has been swallowed, induced vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR Administer two 1/100 grain atropine tablets immediately. NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport patient to the nearest physician.

NOTE TO PHYSICIAN

WARNING SYMPTOMS Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10 minutes intervals in doses of 1 to 2 milligrams, until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural draining. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitor may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

Methyl Parathion Emulsible Co
low-volume ground applicators
commended purposes and at r

INS

For control of aphids, boll
and thrips, apply Methyl Para
0.25 to 0.50 lbs. actual per a

Apply with sufficient water
first appear and repeat appli
not apply within 15 days of
dosage in 5 to 15 gallons of

Mix with water by agitation;
reagitate before use.



KERR-MCGEE CHEMICAL CORP.

GENERAL OFFICES: CHARLOTTE, N.C. OR ANCONA
NEW BERN, NORTH CAROLINA • CHARLESTON, SOUTH CAROLINA

[2342-796]

METHYL PARATHION

Emulsible Concentrate

of Methyl Parathion per gallon

ACTIVE INGREDIENTS:

O,O-Dimethyl O-p-nitrophenyl thiophosphate
Xylene Aromatic Petroleum Solvent

INERT INGREDIENTS:

ACCEPTED

Jun-12-1967

44.30%
50.70%
5.00%
100.00%

USDA Reg. No. 9132-73

ach

ANTIDOTE AND FIRST AID TREATMENT: Atropine is the emergency antidote for Methyl Parathion poisoning. Consult your physician about obtaining an adequate supply of 1,100 grain atropine tablets for emergency use. CALL A PHYSICIAN IMMEDIATELY IN ALL CASES OF SUSPECTED POISONING.

INTERNAL: If the material has been swallowed, induced vomiting immediately. This may be done by introducing a finger into the throat or by giving warm salt water (1 tablespoon of salt to a glass of water). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: If the material has been spilled on the skin, immediately remove patient from the vicinity of the insecticide, remove all contaminated clothing, and wash skin with soap and running water. If the material gets into the eyes, wash immediately with running water for at least ten minutes.

IF WARNING SYMPTOMS APPEAR Administer two 1,100 grain atropine tablets immediately. NEVER ADMINISTER ATROPINE UNLESS WARNING SYMPTOMS APPEAR. Keep patient prone and quiet. Start artificial respiration immediately if patient is not breathing. Transport patient to the nearest physician.

NOTE TO PHYSICIAN

WARNING SYMPTOMS Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10 minutes intervals, in doses of 1 to 2 milligrams, until complete reorganization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitor may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Avoid any further exposure until time for cholinesterase regeneration has been allowed as determined by blood test.

2342-796

GENERAL DIRECTIONS

Methyl Parathion Emulsible Concentrate can be used in conventional hydraulic sprayers, low-volume ground applicators, or airplane sprayers. Use this product only for recommended purposes and at recommended dosages.

INSECT CONTROL ON COTTON

For control of aphids, boll weevil, cotton leaf worm, fleahopper, red spider mites, and thrips, apply Methyl Parathion Emulsible Concentrate at the rate of $\frac{1}{2}$ to 1 pint (0.25 to 0.50 lbs. actual) per acre.

Apply with sufficient water to give uniform coverage. Begin application when insects first appear and repeat application as often as necessary to maintain control. Do not apply within 15 days of hand picking. For application by airplane, mix required dosage in 5 to 15 gallons of water for adequate coverage per acre.

MIXING DIRECTIONS

Mix with water by agitation; agitate the mixture while spraying. If allowed to stand, reagitrate before use.

KERR-MCGEE CHEMICAL CORP.

GENERAL OFFICES: OKLAHOMA CITY, OKLAHOMA
NEW BERN, NORTH CAROLINA • CHARLESTON, SOUTH CAROLINA

