

Feb. 15, 1967 2342-818

THIODAN^R 2-E

Emulsifiable Concentrate
Contains 2 lbs. of endosulfan per gallon

ACTIVE INGREDIENTS:

Endosulfan [®]	24.0%
Xylene type solvent	69.0%
INERT INGREDIENTS:	7.0%
TOTAL	100.0%

*Hexachlorocyclohexane-2,4,3-benzodioxathiepin oxide

^RThiodan is a registered trademark of Farbwerke Hoechst A.G. - U.S. Patent No. 2,799,685.

WARNING (18 Pt. Type)

KEEP OUT OF REACH OF CHILDREN (12 Pt. Type)

Hazardous if swallowed, inhaled or absorbed through skin! Do not breath spray mist. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Wash thoroughly with soap and water after handling, and before eating or smoking; wear clean clothing. During commercial or prolonged exposure in spray-mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type passed by the U. S. Department of Agriculture for THIODAN protection. Workers entering treated areas the day of application should wear protective clothing.

THIS PRODUCT IS POISONOUS TO FISH AND WILDLIFE. KEEP OUT OF ANY BODY OF WATER. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT OF SPRAY FROM AREAS TREATED. RINSE EQUIPMENT AND CONTAINERS AND DISPOSE OF WASTES BY BURYING IN NON-CROP LANDS AWAY FROM WATER SUPPLIES. CONTAINERS SHOULD BE DISPOSED BY PUNCHING HOLES IN THEM AND BURYING WITH WASTES.

CAUTION: Workers entering treated areas the day of application should wear protective clothing.

DIRECTIONS FOR USE

Recommended dosage refer to amount of Thiodan 2-E per acre. Mix with enough water to obtain uniform full coverage. Apply when signs of their damage appear, and repeat at 7-10 day intervals or as necessary.

DIRECTIONS FOR USE (continued)

HEAD LETTUCE: To control aphids, leafhopper, green peach alysiid, flea beetle, black moth larvae, and other insects, use 1 1/2-2 quarts. Do not apply more than 3 applications during growing period. Do not apply within 7 days of harvest. Remove wrapper leaves. Do not use on crop refuse or live plants.

ORNAMENTALS (Greenhouse and outdoors): To control aphids, white fly and cyclamen mite use 1 quart in 100 gallons of water and spray for thorough coverage. In greenhouse applications wear mask or respirator passed by U.S.D.A. for Thiodan protection. Do not use on birch trees. May cause injury on Bonnafon Delume and certain other varieties of chrysanthemums.

PEPPERS: To control aphids and flea beetles use 2 and 1 quarts respectively. Do not apply within 7 days of harvest.

TOMATOES: To control flea beetle, Colorado potato beetle, leafhopper and aphids, use 1 quart and apply as needed up to harvest.

STRAWBERRIES: To control strawberry aphid and meadow spittlebug use 1 1/2 quarts. Do not apply within 7 days of picking. Do not apply more than twice during any 30 day period.

LEGGOS: To control flea beetle, hornworm, budworm and grasshopper use 1-2 quarts. Do not apply within 7 days of harvest. Use seed treatment 1 quart in 100 gallons of water. Apply 1 quart per 1000 square yards.

TOMATOES & PEPPERS: To control blister beetle, Colorado potato beetle, flea beetle, hornworm, green alfalfa and white fly, use 1 quart per acre. Do not apply within 7 days of harvest.

WARNINGS - PLANTS

KEEP AWAY FROM LEAVES AND FRUIT OF GREENHOUSE PLANTS. DO NOT APPLY TO PEACHES, BERRIES OR FRUIT WHICH WILL BE BLOWN OFF. DO NOT USE ON PLANTS IN BLOSSOM.

Thiodan is a registered trademark of Farbwerke Hoechst A.G. - U.S. Patent No. 2,799,685.

WARNING (18 Pt. Type)

KEEP OUT OF REACH OF CHILDREN (12 Pt. Type)

Hazardous if swallowed, inhaled or absorbed through skin! Do not breath spray mist. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing, and flush skin or eyes with plenty of water; for eyes, get medical attention. Wash thoroughly with soap and water after handling, and before eating or smoking; wear clean clothing. During commercial or prolonged exposure in spray-mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type passed by the U. S. Department of Agriculture for THIODAN protection. Workers entering treated areas the day of application should wear protective clothing.

THIS PRODUCT IS POISONOUS TO FISH AND WILDLIFE. KEEP OUT OF ANY BODY OF WATER. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT OF SPRAY FROM AREAS TREATED. RINSE EQUIPMENT AND CONTAINERS AND DISPOSE OF WASTES BY BURYING IN NON-CROP LANDS AWAY FROM WATER SUPPLIES. CONTAINERS SHOULD BE DISPOSED BY PUNCHING HOLES IN THEM AND BURYING WITH WASTES.

CAUTION: Workers entering treated areas the day of application should wear protective clothing.

DIRECTIONS FOR USE

Recommended dosages refer to amount of Thiodan 2-3 p.p.a. Mix with enough water to obtain uniform full coverage. Apply when insects or their damage appear, and repeat at 7-10 day intervals or as necessary.

CABBAGE, CAULIFLOWER, BRASSICA: To control cabbage aphid, cabbage looper, imported cabbage worm, diamond-back moth larvae, green-striped cabbageworm, flea beetle, meadow spittlebug, tinker bug. Use 1-2 quarts. Do not apply to cauliflower after edible parts begin to form. Do not apply to Brussels sprouts and cabbage within 7 days of harvest.

CUCUMBERS, MELONS, PEPPERS, SWEET SQUASH, SWEET POTATOS: To control aphids, cucumber beetle, striped cucumber beetle, squash vine borer, melonworm, flower bud, stem and vine borer. Use 1-2 quarts. Application may be repeated as needed.

ORNAMENTALS (Greenhouse and outdoors): To control aphids, white fly and cyclamen mite use 1 quart in 100 gallons of water and spray for thorough coverage. In greenhouse applications wear mask or respirator passed by U.S.D.A. for Thiodan protection. Do not use on birch trees. May cause injury on Dianthus barbatus and certain other varieties of chrysanthemums.

PEPPERS: To control aphids and flea beetles use 2 and 1 quarts respectively. Do not apply within 1 day of harvest.

POTATOES: To control flea beetle, Colorado potato beetle, leafhopper and aphids, use 1 quart and apply as needed up to harvest.

STRAWBERRIES: To control strawberry aphid and meadow spittlebug, use 1-2 quarts. Do not apply within 4 days of picking. Do not apply at intervals of less than 15 days; or more than twice during any 35 day period.

TOBACCO: To control flea beetle, hornworm, budworm and green stink bug, use 1-2 quarts. Do not apply within 5 days of harvest. In seed beds use 1 quart in 100 gallons of water and apply 6 gallons of water per 100 square yards.

TOMATOES & EGGPLANTS: To control blister beetle, Colorado potato beetle, flea beetle, hornworm, green stink bug and white fly, use 1 quart per acre. Do not apply within 1 day of harvest.

WARNING! PLANTS!

KEEP AWAY FROM NEARBY PLANTS. KEEP CONTAINER CLOSED. CONTAINERS SHOULD BE KEPT IN A SAFE PLACE. DO NOT STORE IN A PLACE WHERE CHILDREN CAN GET AT THEM.

CONDITIONS OF SALE

The seller warrants that the product to be true to the label and that, it makes no representation or warranty, and is not responsible for any damage or loss from the use or handling of the material, whether or not such use or handling is in accordance with directions or instructions on the label or user acceptable to the above conditions.

Manufactured by
Franklin Chemical Co.
Oklahoma City, Oklahoma
U.S.D.A. Registration No. 933-100



Contains 2 1/2 lbs. of active ingredient

Net Weight	2.500
Net Content	0.750
Net Weight	1.750
TOTAL	1.750

Hexachlorocyclopentadiene-2,4,6-trichloro-1,3-dioxane

Infidan is a registered trademark of Farbwerke Hoechst A.G. - U.S. Patent No. 2,799,685.

WARNING (15 Fl. Oz.)

KEEP OUT OF REACH OF CHILDREN (15 Fl. Oz.)

Hazardous if swallowed, inhaled or absorbed through skin! Do not breathe spray. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Wash thoroughly with soap and water after handling, and before eating or smoking; wear clean clothing. During commercial or prolonged exposure, in spray-mixing and loading operations, wear clean synthetic rubber gloves and a mask or respirator of a type passed by the U. S. Department of Agriculture for TOXIC agents. Workers entering treated areas should use appropriate protective clothing.

WILDLIFE. KEEP OUT OF REACH OF WILDLIFE.

USE DIRECTIONS: For control of leaf beetles, beetles, grasshoppers, crickets, and locusts, apply 1 quart per acre. Do not apply within 10 days of harvest. For control of grasshoppers, crickets, and locusts, apply 1 quart per acre. Do not apply within 10 days of harvest. For control of grasshoppers, crickets, and locusts, apply 1 quart per acre. Do not apply within 10 days of harvest.

ACCEPTED

FEB 15 1967

UNDER THE PROVISIONS OF THE FEDERAL FOOD, DRUG, AND COSMETIC ACT FOR LISTING OF PESTICIDES REGISTERED IN THE UNITED STATES