

ACCEPTED
JAN 3 1967
UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE OF AN ...
FOR EQUIV ...
ED UNDER NO. 91373 SUBJECT
TO ATTACHED COMMENTS.

2342-855

FRONT PANEL COPY

Net Contents - 8 ounces
Makes 25 gallons of spray.

GARDEN & LAWN
FUNGICIDE
Contains 50% Captan

Controls: Black Spot on roses, Brown Patch
on lawns, Damp-off on bulbs & cuttings, and
many diseases on fruits and vegetables.

CAUTION: KEEP OUT OF REACH OF CHILDREN!
See side panel for additional required cau-
tion statements.

SIDE PANEL COPY

1-3-67
9132-173

2342-835

CAUTION

Avoid inhalation of dust or spray mist.
Avoid prolonged or repeated contact with skin. Wash hands and face after use.
Avoid contamination of feed and foodstuffs.
Dispose of container promptly when empty.

ACTIVE INGREDIENT	By Wt.
Captan*	50%
INERT INGREDIENTS	50%
TOTAL	<u>100%</u>

*N/ (trichloromethyl) thio/-4-cyclohexene-1, 2-dicarboximide.

NOTE:
This can contains 8 ounces of material as indicated. Considerable settling may have occurred due to shipping and handling.


1-3-67
9132-173

2342-835

SIDE PANEL COPY

CONDITIONS OF SALE

The seller guarantees the material sold to be true to the label. Further than that, it makes no representation or warranty, and it shall not be held responsible for any injury to crop, soil, person, or property, resulting from the use or handling of said material, whether used in accordance with directions on the label. The buyer and/or user accept this material subject to the above conditions of sale.



2342-835

BACK PANEL COPY

GARDEN & LAWN FUNGICIDE

DIRECTIONS: Use 8 teaspoonfuls per gallon of water unless otherwise specified. Stir well before using and agitate frequently during application.

APPLE TREES:

CONTROLS: Apple scab, fruit spot, bitter rot. **APPLICATION:** Apply in pre-blossom, petal fall and cover sprays.

PEACHES AND CHERRIES:

CONTROLS: Brown rot and cherry leaf spot. **APPLICATION:** Apply at pink bud, full bloom and petal fall stages. Repeat at 14 day intervals after petal fall.

TOMATO AND POTATO:

CONTROLS: Early and late blights, anthracnose and gray leafspot of tomatoes. **APPLICATION:** Use 16 teaspoonfuls per gallon water. Apply at first sign of infection and repeat at 7 day intervals to maintain control.

ROSES:

CONTROLS: Black Spot. **APPLICATION:** Use 3 tablespoonfuls per gallon water. Apply at first sign of disease. If powdery mildew becomes a problem switch to Folpet fungicide.

TUBEROUS BEGONIAS:

CONTROLS: Powdery Mildew. **APPLICATION:** Use 2 tablespoonfuls per gallon water. Apply at first sign of disease and repeat at 10 day intervals.

BULBS (glads and begonias):

CONTROLS: Rot and damp-off. **APPLICATION:** Use 8 tablespoonfuls per gallon water. Dip bulbs for 20-30 minutes, drain and plant.

CUTTINGS (azaleas, carnations, chrysanthemums):

CONTROLS: Damp-off. **APPLICATION:** Use 5 tablespoonfuls per gallon water. Dip cuttings before bedding.

Cont'd.

Cont'd.

LAWNS:

CONTROLS: Brown patch, leaf spot, damp-off, root rot, Helminthosporium (melting-out). **APPLICATION:** Apply at rate of 1 gallon per 100 square feet at first sign of disease. Make 3 applications 7 days apart to maintain control.

1-3-67
9132-173
THE F. B. I.
SEARCHED
SERIALIZED
INDEXED
FILED

2342-835