

[2342-841]

1% ROTENONE DUST

ACTIVE INGREDIENTS:

Rotenone	1.00%
Other Carb Resins	2.00%

INERT INGREDIENTS:

97.00%

TOTAL

100.00%

CAUTION (18 Point Type)

KEEP OUT OF REACH OF CHILDREN (12 Point Type)

CONDITIONS OF SALE

This material may be irritating to some individuals.
Use reasonable care to avoid inhaling it.

THIS PRODUCT IS POISONOUS TO FISH.
KEEP OUT OF ANY BODY OF WATER.

DIRECTIONS FOR USE

These are general suggestions for insect control.
Growers should confer with Federal and State
authorities for recommendations best suited to
their local conditions.

PEAS, for pea aphid.

CABBAGE, CAULIFLOWER & BROCCOLI, for flea
beetles.

BEANS, for bean beetles.

ASPARAGUS, for asparagus beetles on market shoots.

PEACHES, GRAPES, for Japanese beetles.

STRAWBERRIES, for spittle bug.

RASPBERRIES, for raspberry fruit worm.

CURRENTS AND GOOSEBERRIES, for gooseberry fruit worm.

CHERRIES, for fruit fly.

BLUEBERRIES, for blueberry maggot.

SWEET CORN, for European Corn Borer. Make 4
applications every 4 to 5 days from the time

the eggs are about to hatch.

Dust thoroughly and uniformly, both upper and
lower surfaces of leaves, at rate of 25 to 50
pounds to the acre, depending upon size of plants.

Apply when insects appear and continue at weekly
intervals throughout the growing season. Do not
apply within 1 day of harvest.

The seller guarantees the material
sold to be true to the label. Further
than that, it makes no representation
or warranty, and it shall not be
held responsible for any injury to
crop, soil, person or property,
resulting from the use or handling
of said material, whether used in
accordance with directions or not.
The buyer and/or user accept this
material subject to the above
conditions of sale.

Manufactured By
KERR-McGEE CHEMICAL CORP.
Oklahoma City, Okla.

U.S.D.A. Registration No. 9132-