

PARATHION 8-E

Contains 1 lb. Parathion per gallon

ACTIVE INGREDIENT


Parathion*	81.5%
INERT INGREDIENTS	18.5%
TOTAL	100.0%

*O,O-Diethyl O-p-nitrophenyl thiophosphate

DANGER (18 Point Type)

KEEP OUT OF REACH OF CHILDREN (12 Point Type)

Poisonous if swallowed, inhaled or absorbed through skin! Rapidly absorbed through skin! Do not get in eyes, on skin, or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated area should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and food-stuffs. Wash hands, arms and face thoroughly with soap and water before eating or drinking. Wash all contaminated clothing with soap and hot water before re-use.



ANTIDOTES: If swallowed - give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. Call a physician immediately.

If on skin - In case of contact remove contaminated clothing and immediately wash skin with soap and water.

NOTE: PHYSICIAN: In case of poisoning, give atropine, preferably by injection, grains 1/32 to 1/16 (2 to 4 mg.) at once and every hour or until needed to achieve and maintain full atropinization. Wear gloves and gown. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continually for 48 hours.

DIRECTIONS FOR USE

Crabapple - To control European Fruit Tree and Sparganathis Fruit Tree, use 1 pt. per acre in sufficient water to insure thorough coverage. Do not apply within 30 days of harvest.

Potato - To control Colorado Potato Beetle, Flea Beetle, Leaf Miner and Serpentine Leaf Miner, use 1 pt. per acre in enough water to insure thorough coverage. Do not apply to tubers within 1 day of harvest or to tubers or tubers within 30 days of harvest.

Tobacco - To control White and Flea Beetles use 1 pt. per acre (for ground application) in enough water to insure thorough coverage. For air application use 1/3 pt. per acre. Do not apply within 10 days of harvest or 10 days of cutting. Avoid drift to other plants, especially other tobacco plants.

PARATHION 8-E

-2-

CAUTION

This product will kill fish and wildlife. Keep out of any body of water.

Apply this product only as specified on this label.

Do not apply when weather conditions favor drift of spray from areas treated.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

CONDITIONS OF SALE

The seller guarantees the material sold to be true to the label. Further than that, it makes no representation or warranty, and it shall not be held responsible for any injury to crop, soil, person or property, resulting from the use or handling of said material, whether used in accordance with directions or not. The buyer and/or user accept this material subject to the above conditions of sale.

Manufactured by
KERR-McGEE CHEMICAL CORP.
Oklahoma City, Okla.

U.S.D.A. Registration No. 9132-