

MANEB THIODAN^R DUST

ACTIVE INGREDIENTS:

Maneb (Manganese ethylenebisdithio- carbamate)	5.60%
Endosulfan*	3.00%
INERT INGREDIENTS:	91.40%
TOTAL	100.00%

*Hexachlorohexahydromethano-2,4,3-
benzodioxathiepin oxide.

^RThiodan is a registered trademark of
Farbwerke Hoechst A.G. U. S. Patent
No. 2,799,685.

CAUTION (18 Point Type)
KEEP OUT OF REACH OF CHILDREN (12 Point Type)

armful if swallowed, inhaled or absorbed
through skin. Avoid breathing dust or spray
mist. Avoid contamination of feed and food-
stuffs. Avoid contact with skin, eyes and
clothing. In case of contact with skin or
eyes flush with plenty of water; for eyes,
get medical attention. Wash with soap and
water after handling and before eating or
smoking; wear clean clothing. Workers entering
treated areas the day of application should
wear protective clothing. Do not apply or
allow to drift to acres occupied by unprotected
humans or beneficial animals. Toxic to fish
and wildlife, do not contaminate streams, lakes
or ponds with this material. Apply this product
only as specified on this label. Dispose of
astes by burying in non-crop lands away from
water supplies. Containers should be disposed
by burying with wastes, or by burning (Keep out
of smoke).

DIRECTIONS FOR USE

Follow application rates given in the following
directions. Dust to give a full, uniform cover-
age to the particular crop. Apply when diseases
or insects first appear. Repeat at 7-10 day in-
tervals, or as necessary.

POTATOES: To control Flea Beetle, Colorado
Potato Beetle, Leafhopper, Aphids and Early
and Late Blight, use 30-35 lbs. per acre. Apply
as needed up to harvest.

TOMATOES: To control Aphids, Blister Beetle,
Colorado Potato Beetle, Flea Beetle, Green
Stink Bug, Hornworm, White Fly, Anthracnose,
Early Blight, Late Blight and Septoria Leaf
Spot, use 20-30 lbs. per acre. Do not apply
within 1 day of harvest.

DIRECTIONS FOR USE (Continued)

CUCUMBERS, MELONS, PUMPKINS, SUMMER
SQUASH & WINTER SQUASH: To control
Aphids, Cucumber Beetle, Striped
Cabbage Flea Beetle, Squash Vine Borer,
Downey Mildew, Anthracnose and
Alternaria, Leaf Spot, use 30-35 lbs.
per acre. Do not use Maneb on young
melons before runners form. For Vine
Borers apply weekly to flower buds,
stems and vines when moths first
appear. Application may be made up
to the day of harvest.

CONDITIONS OF SALE

The seller guarantees the material
sold to be true to the label.
Further than that, it makes no
representation or warranty, and it
shall not be held responsible for any
injury to crop, soil, person or
property, resulting from the use or
handling of said material, whether
used in accordance with directions
or not. The buyer and/or user accept
this material subject to the above
conditions of sale.

Manufactured By
KERR-McGEE CHEMICAL CORP.
Oklahoma City, Okla.

U.S.D.A. Registration No. 9132-

4132 209

CO

2-342-853