

NET WEIGHT 2-1/2 LBS.

DICHLONE 50-W

ACTIVE INGREDIENTS:

Dichlone (2,3-dichloro-1,4-Naphthoquinone)..... 50.0%

INERT INGREDIENTS:..... 50.0%

TOTAL 100.0%

CAUTION (18 Point Type)

KEEP OUT OF REACH OF CHILDREN (12 Point Type)

May cause skin irritation. Care should be taken in handling and applying this material. Wash with soap and water after using and before eating or smoking. Avoid inhalation of dust. Avoid contact with eyes, skin or on clothing. Wear protective clothing. Avoid storage near feed and food products.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by burning. (Keep Out Of Smoke).

THIS PRODUCT IS TOXIC TO FISH
KEEP OUT OF LAKES, STREAMS, AND PONDS.

Apply this product only as specified on this label.

DIRECTIONS FOR USE: (two level teaspoonfuls per gallon equals 1 lb. per 100 gallons).

Apple Scab - use 1/4 lbs. per 100 gallons of water on protective schedule at 5-7 day intervals from delayed dormant through the early cover sprays. Use 1/2 lb. per 100 gallons in applications made up of 48 hours after the start of a wetting period for post infection protection.

Peach Brown Rot - for blossom and twig rot blossom blight use 1/2 lb. per 100 gallons of water in pink bud stage, full bloom and just before shuck fall. When cool wet weather occurs, resulting in a long bloom period, apply Phygon every 4 days. As a pre-harvest spray use 1/2 lb. per 100 gallons at 14 and 7 days before harvest.

Peach Leaf Curl - use 1 lb. per 100 gallons of water as a dormant spray just before the buds begin to swell.

Cherry Brown Rot Blossom Blight - use 1/2 lb. per 100 gallons of water at popcorn, full bloom and petal fall stages.

Tomato Leaf Blights - late blight and gray mold (botrytis) - use 3/4 lb. per 100 gallons of water at 5-7 day intervals, beginning when plants are 4" high. To control the same diseases in tomato plant beds, use 1 lb. per 100 gallons of water at 5-7 day intervals beginning when the plants emerge from the ground.

ACCEPTED
MAY 22 1967
UNDER THE FEDERAL
FUNGICIDE AND FUNGICIDE ACT
FOR ECONOMIC INSECTICIDE REGISTERED
ED UNDER NO. 11221-100 SUBJECT
TO ATTACHED COMMENTS

2-42-874

DIRECTIONS FOR USE (Continued)

Celery Blights - use 1/2 lb. per 100 gallons of water every 5-7 days during infection period. Use 200 gallons of dilute spray per acre. Make 3-4 applications about 7-10 days after plants are set.

Azalea Petal Blight - use 1/2 lb. per 100 gallon of water every 5-7 days as the blossoms show color and open.

Rose Black Spot - use 1/4 lb. per 100 gallons of water every 5-7 days. Begin applications early before disease is apparent. Stop applications when signs of infection.

Potato Late Blight - use 1/2 lb. Phygon per 100 gallons of water every 5-7 days after blossom period. If blight threatens prior to this time, use a Maneb, Nabam or Copper Fungicide.

NOTE: Do not use Phygon with Bordeaux Mixture, Lime, Sulfur, Organic Mercuries, Oils or Oil Emulsions.

CONDITIONS OF SALE

The seller guarantees the material sold to be true to the label. Further than that, it makes no representation or warranty, and it shall not be held responsible for any injury to crop, soil, person or property, resulting from the use or handling of said materials, whether used in accordance with directions or not. The buyer and or user accept this material subject to the above conditions of sale.

Manufactured By
Kerr-McGee Chemical Corp.
Oklahoma City, Okla.

U.S.D.A. Registration No. 9132-