


CHLORDANE 8-E

Contains 8 lbs. Chlordane Per Gallon



ACTIVE INGREDIENTS:

Technical Chlordane*..... 73%
Petroleum distillates..... 22%

INERT INGREDIENTS:..... 5%
TOTAL..... 100%

*Equivalent to 43.7% Octachloro-4,7-methanotetrahydroindane and 29.2% related compounds

CAUTION (18 Point Type)

KEEP OUT OF REACH OF CHILDREN (12 Point Type)

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin wash with soap and water. Avoid contamination of feed and foodstuffs. Do not allow children or pets on treated lawns until Chlordane has been washed into the soil and lawn is dry.

DIRECTIONS FOR USE:

SUBTERRANEAN TERMITES: A chemical barrier between the termites in the soil and wooden structures and foundations is necessary to obtain good control. For buildings with deep foundations dig a trench 30 inches deep and 10 inches wide next to the foundation wall. For buildings with shallow foundations do not dig below the top of the footing. Add 1 pint of Chlordane 8-E to 12 gallons of water and apply in the trench at the rate of 4 gallons per 5 linear feet. Apply 1 gallon of the mixture per 10-square feet as an overall treatment under concrete slabs and under attached porches.

LAWN AND ORNAMENTAL TURF: Ants, army worms, crickets, grasshoppers, mosquito and spiders: Use 1 pint of Chlorodane 8-E in enough water to thoroughly spray one acre of lawn including ornamental plants. Note: Do not direct foliage sprays onto feed and forage crops or crops for human consumption.

Chiggers and ticks: Use 1 1/2 pints Chlordane 8-E in enough water to thoroughly spray one acre of lawn

Chinch bugs, lawn moths and sod webworms: Use 3 pints of Chlordane 8-E in enough water to thoroughly spray one acre of lawn or turf.

Japanese beetle grubs, white grubs and white fringed beetle larvae: Use 2 quarts of Chlordane 8-E in 100 gals. of water to thoroughly spray one acre of lawn or turf. Wash Chlordane into the soil immediately after application.

AGRICULTURAL APPLICATIONS:

Wireworms in beans, beets, celery, cole crops, (cabbage, broccoli, brussel sprouts, cauliflower, collards, kale & kohlrabi), corn, onions, peppers, potatoes, tobacco and turnips: Use 1 1/2-2 1/2 quarts of Chlordane 8-E in enough water to thoroughly spray one acre of soil before planting. Work into the top few inches of the soil before planting.

Root maggots on cole crops, (cabbage, broccoli, brussel sprouts, cauliflower, collards, kale & kohlrabi), radishes and turnips: Apply as a preplant treatment or at planting time only. Use 1 quart of Chlordane 8-E in enough water to thoroughly spray one acre of soil. Work Chlordane into the top few inches of the soil.

Flea beetle larvae, Japanese beetle grubs and white grubs in potato and tobacco fields: Use 1 1/2-2 1/2 quarts of Chlordane 8-E in enough water to thoroughly spray one acre of soil before planting or transplanting. Work into the top few inches of the soil before planting.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

CONDITIONS OF SALE: The Kerr-McGee Chemical Corp. and the seller make no warranty of any kind, expressed or implied on the use of this material and assume no responsibility whatsoever for any personal injury, property damage, or other type of loss resulting from the use or handling of this material. The user assumes all risks for use or handling other than in accordance with directions.

NET CONTENTS: 1 Gal. 5 Gals.
30 Gals. 55 Gals.

Manufactured By:
KERR-McGEE CHEMICAL CORP.
Kerr-McGee Building
Oklahoma City, Oklahoma

U.S.D.A. REG. NUMBER 9132-