

Available Phosphoric Acid
(P₂O₅).....16.00%
Soluble Potash (K₂O).....16.00%

DIRECTIONS FOR USE

DI-SYSTON is an effective systemic insecticide for early season crop protection which has been combined with fertilizer for efficient and uniform application. For best results apply at time of planting. Do not use on other crops used for food or forage. Use only according to label directions. Applications at rates higher than those shown on label may result in illegal crop residues.

POTATOES: For control of Aphids, Flea beetles (except Tuber flea beetles), Leafhoppers, Potato psyllid, Southern potato wireworm and Colorado potato beetle (early season reduction).

RATE: Apply 1350 to 2000 lbs. (2 to 3 lbs. active insecticide) per acre. Apply specific dosage per acre in a band on each side of the seed furrow at time of planting only. Applications may be made as a side-dressing after plants have become established. Do not apply within 75 days of harvest on potatoes. Use the higher rate in organic soils.

Note: Use either at planting time or as a side-dressing but not both treatments.

(R) DI-SYSTON is the Registered Trademark of Farbenfabriken Bayer A-G, Chemagro Corporation, licensee. U.S.D.A. REGISTRATION NO. 9132-X

Sulfate. Atropine Sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.


HIGHLY TOXIC TO FISH AND WILDLIFE. DO NOT CONTAMINATE STREAM, WOODLAND AND OTHER NON-CROP AREAS. CONTAINERS SHOULD BE DISPOSED OF BY BURYING OR BY BURNING. (KEEP OUT OF SMOKE)

NOTICE

The Kerr-McGee Chemical Corporation and the seller make no warranty of any kind, express or implied on the use of this material and assume no responsibility whatsoever for any personal injury, property damage or other type of loss resulting from the use or handling of this material. The user assumes all risks for use or handling whether in accordance with directions or not.

The principle active ingredient of this product is patented under U.S. Patent No. 2,759,010 of Chemagro Corporation, Kansas City, Missouri, and the purchase of this product includes a limited license under said patent to use or reformulate the product only in the manner provided by the label. The purchaser obligates itself to notify Chemagro Corporation of any proposed use or reformulation of such licensed product not provided by said recommended applications, and to obtain Chemagro's approval in writing before making any such use or reformulation of the licensed product. Any use, or reformulation intended for use of the licensed product in the home and garden field is specifically prohibited by said license

Manufactured by
Kerr-McGee Chemical Corporation
Oklahoma City, Oklahoma



DECEMBER 29, 1967

(K-M SEAL)

NET WEIGHT 80 LBS.

8-16-24 + D for Tobacco
Contains 0.50% *DI-SYSTON[®] or 10
pounds active per ton
For Use Under Tobacco

STATEMENT OF INGREDIENTS FOR INSECT
CONTROL

ACTIVE INGREDIENTS:

*O,O-Diethyl S- 2-(ethylthio)
ethyl phosphorodithioate... 0.50%
INERT INGREDIENTS..... 99.50%
TOTAL.....100.00%

GUARANTEED ANALYSIS FOR PLANT FOOD

Total Nitrogen (N)..... 8.00%
Available Phosphoric Acid
(P₂O₅).....16.00%
Soluble Potash (K₂O).....24.00%

DIRECTIONS FOR USE

DI-SYSTON is an effective systemic insecticide for early season crop protection which has been combined with fertilizer for efficient and uniform application. For best results apply at time of planting. Do not use on other crops used for food or forage. Use only according to label directions. Applications at rates higher than those shown on label may result in illegal crop residues.

TOBACCO: For control of Rootworms, Aphids, Mites, Leafhoppers, Thrips, Flea beetle and Wireworm.

Rate: Broadcast 700 to 800 lbs. (3½ to 4 lbs.) of active Di-Syston per acre. Apply specific rate per acre after plowing and disc thoroughly into the soil surface to a depth of 4 to 6 inches before planting tobacco. Do not apply more than 800 lbs. per acre.

OR

This product may also be applied at time of transplanting as a furrow treatment. Apply 700 to 800 lbs. per acre on one side or on each side of the tobacco plant at time of transplanting. Do not apply fertilizer in direct contact with the roots of the tobacco transplant due to the danger from injury. For best placement this product should be placed 3 inches to the side or sides of the transplant and one inch deeper than the root system. Do not apply more

WARNING

KEEP OUT OF REACH OF CHILDREN

Poisoning may occur as a result of excessive exposure. May be fatal if swallowed. May be absorbed through skin. Do not breathe dust. Avoid contact with skin, eyes, or clothing. Do not contaminate feed or food. Wash thoroughly after handling.

Antidote: Atropine sulphate.
TO PHYSICIAN: Atropine Sulfate is antidotal, 2-PAM is also antidotal and may be administered in conjunction with atropine.

HIGHLY TOXIC TO FISH AND WILDLIFE. TO PROTECT WATER SUPPLIES, FISH AND WILDLIFE, DO NOT CONTAMINATE ANY BODY OF WATER NOR APPLY TO ANY AREA NOT SPECIFIED ON THE LABEL.

NOTICE

The principle active ingredient of this product is patented under U.S. Patent No. 2,759,010 of Chemagro Corp., Kansas City, Missouri, and the purchase of this product includes a limited license under said patent to use or reformulate the product only in the manner provided by the recommended applications on the present label. The purchaser obligates itself to notify Chemagro Corp. of any proposed use or reformulation of such licensed product not provided by said recommended applications and to obtain Chemagro's approval in writing before making any such use or reformulation of the licensed product. Any use, or reformulation intended for use, of the licensed product in the home or garden field is specifically prohibited by