

[2342-913]



Net Contents
50 Pounds

DY-BAIT CROP INSECTICIDE

DIRECTIONS FOR USE

Apply DY-BAIT at specified rate per acre for the control of surface-feeding cutworms, salt marsh caterpillars, armyworms and crickets.

Applications may be made in a band over planted row or around the stems of individual plants. If such methods are not practical, apply the bait as a broadcast treatment over the infested area.

If field has a history of previous cutworm damage or if cutworms are suspected, apply DY-BAIT before or soon after planting or transplanting.

RECOMMENDED DY-BAIT APPLICATIONS

CROP	POUNDS DY-BAIT/ACRE	REMARKS
FIELD CROPS		
Alfalfa	20	Do not apply more than one application per cutting nor within 14 days of cutting or grazing.
Barley	20	Do not apply within 21 days of harvest. Do not graze treated fields.
Flax		
Oats		
Wheat		
Corn	20	Apply once per season when plants are 3-12 inches high. Do not use corn for food, feed, or ensilage within 40 days of treatment.
Cotton	30	Do not apply within 7 days of picking. Do not graze livestock in treated fields within 14 days after application.

Contains 5% [®] DYLOX

ACTIVE INGREDIENT	BY WT.
Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate†	5%
INERT INGREDIENTS	95%
TOTAL — 100%	

†U. S. Patent No. 2,701,225 Canadian Patent No. 529,379
EPA Reg. No. 9132-290

WARNING: Keep Out of the Reach of Children. See additional cautions on this label.

STOP: Read the Label before use.

VEGETABLES	
Beans (Dry & Snap)	30
Beans (Lima)	30

Do not apply within 14 days of harvest. Do not feed treated vines to meat or dairy animals.

Do not apply more than twice per season nor harvest for food, feed or forage within 20 days of treatment.

DANG

This product is toxic. Treated areas may be hazardous. Conditions favor drift. Run-off is likely to occur. Avoid contact with equipment, or

Do not use on crops. Use only according to directions above those shown. Do not treat food crops grown

Do not re-use the container and dispose of waste water supplies. Dispose of wastes or by burning

May be harmful if inhaled. Avoid contact with skin. Avoid contact with eyes. Wash after handling. Wash with hot water before re-dressing. Do not contaminate clothing. Not for use or if poisoning occurs,

ACCEPTED

JAN 13 1972

9132-290

DIRECTIONS FOR USE

Apply DY-BAIT at specified rate per acre for the control of surface-feeding cutworms, salt marsh caterpillars, armyworms and crickets.

Applications may be made in a band over planted row or around the stems of individual plants. If such methods are not practical, apply the bait as a broadcast treatment over the infested area.

If field has a history of previous cutworm damage or if cutworms are suspected, apply DY-BAIT before or soon after planting or transplanting.

RECOMMENDED DY-BAIT APPLICATIONS

CROP	POUNDS DY-BAIT/ACRE	REMARKS
FIELD CROPS		
Alfalfa	20	Do not apply more than one application per cutting nor within 14 days of cutting or grazing.
Barley Flax Oats Wheat	20	Do not apply within 21 days of harvest. Do not graze treated fields.
Corn	20	Apply once per season when plants are 3-12 inches high. Do not use corn for food, feed, or ensilage within 40 days of treatment.
Cotton	30	Do not apply within 7 days of picking. Do not graze livestock in treated fields within 14 days after application.
Peanuts	20	Applications may be made in a band over the row or broadcast by air or ground equipment. Three applications may be made before digging. One additional application may be made between digging and harvest.
Safflower	30	Do not apply after plants begin to bloom. Do not apply more than twice per year.
Sugar Beets	30	Do not apply within 28 days of harvest if tops are to be fed to livestock nor within 14 days of harvest if tops are not to be fed.
Tobacco	20	Do not apply after plants are 2 feet tall.
SEED FIELD CROPS		
Soybeans	30	Do not pasture or use treated crop for feed, food or forage.

Contains 5% [®] DYLOX

ACTIVE INGREDIENT	BY WT.
Dimethyl (2,2,2-trichloro-1-hydroxyethyl) phosphonate†	5%
INERT INGREDIENTS	95%
TOTAL — 100%	

†U. S. Patent No. 2,701,225 Canadian Patent No. 529,379
EPA Reg. No. 9132-290

WARNING: Keep Out of the Reach of Children. See additional cautions on this label.

STOP: Read the Label before use.

VEGETABLES		
Beans (Dry & Snap)	30	Do not apply within 14 days of harvest. Do not feed treated vines to meat or dairy animals.
Beans (Lima)	30	Do not apply more than twice per season nor harvest for food, feed or forage within 30 days of treatment.
Cowpeas (Southern peas, blackeyed peas, crowder peas)	30	Do not apply more than twice per season nor within 14 days of harvest. Do not feed treated vines to meat or dairy animals.
Brussels Sprouts Cabbage Cauliflower	20	Do not apply within 21 days of harvest.
Collards Lettuce	20	Do not apply within 28 days of harvest.
Peppers	20	Do not apply within 21 days of harvest.
Pumpkins	20	Do not apply more than once per season nor within 14 days of harvest.
Table Beets	30	Do not apply within 28 days of harvest. Do not eat tops.
Tomatoes	20	Do not apply within 28 days of harvest. Do not graze treated fields.

DANG

This product is toxic. Treated areas may be hazardous. Conditions favor drift. Run-off is likely to occur. Cleaning of equipment, etc.

Do not use on crops. Use only according to label. Above those shown, do not treat food crops grown for human consumption.

Do not re-use the container and dispose of wastes in water supplies. Dispose of wastes or by burning.

May be harmful if inhaled. Avoid contact with skin. Avoid contact with eyes. Wash after handling. Wash hands with hot water before rehandling. Do not contaminate clothing. Not for use on food. If poisoning occurs, medical posure will result in death. **NOTE TO PHYSICIAN:** Also antidotal and atropine.

All statements concerning this product when used as directed. The manufacturer makes no representation concerning this product. Description on the label. DYLOX is Reg. TM of Bayer Benfabriken Bayer AG, Leverkusen, Germany.

DRAFT C

ACCEPTED
 Jan 13, 1972
 [2942-913]
 9132-290
 TO APPLICATOR

Manufactured by **KERR-MCGEE CHEMICAL CORP.** Jacksonville, Florida,

ASCO

OP INSECTICIDE

Product No.
20-730211

ins 5% [®] DYLOX

NT	BY WT.
-trichloro-1-hydroxyethyl)	
	5%
TS	95%
TOTAL — 100%	

01,225 Canadian Patent No. 529,379
A Reg. No. 9132-290

**G: Keep Out of the
dren. See additional cau-
label.**

ad the Label before use.

DANGER TO FISH AND WILDLIFE

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where run-off is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.

RESTRICTIONS

Do not use on crops other than those specified on this label. Use only according to label directions. Applications at rates above those shown may result in illegal crop residues. Do not treat food crops grown in greenhouses.

CONTAINER DISPOSAL

Do not re-use the containers. Rinse equipment and containers and dispose of wastes by burying in non-crop land away from water supplies. Dispose of empty container by burying with wastes or by burning. (Keep out of smoke and fumes).

CAUTION

May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes, and clothing. Do not breathe dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use.

Do not contaminate feed or food. Keep out of reach of children. Not for use or storage in or around the home. If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.

NOTE TO PHYSICIAN: Atropine Sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

CONDITIONS OF SALE

Do not apply within 14 days of harvest. Do not feed treated vines to meat or dairy animals.

Do not apply more than twice per season nor harvest for food, feed or forage within 30 days of treatment.

Do not apply more than twice per season nor within 14 days of harvest. Do not feed treated vines to meat or dairy animals.

Contains 5% [®] DYLOX

INGREDIENT	BY WT.
2,2,2-trichloro-1-hydroxyethyl phosphate	5%
OTHER INGREDIENTS	95%
TOTAL — 100%	

No. 2,701,225 Canadian Patent No. 529,379
EPA Reg. No. 9132-290

**WARNING: Keep Out of the
Children. See additional cau-
tions on this label.**

Read the Label before use.

30	Do not apply within 14 days of harvest. Do not feed treated vines to meat or dairy animals.
30	Do not apply more than twice per season nor harvest for food, feed or forage within 30 days of treatment.
30	Do not apply more than twice per season nor within 14 days of harvest. Do not feed treated vines to meat or dairy animals.
20	Do not apply within 21 days of harvest.
20	Do not apply within 28 days of harvest.
20	Do not apply within 21 days of harvest.
20	Do not apply more than once per season nor within 14 days of harvest.
30	Do not apply within 28 days of harvest. Do not eat tops.
20	Do not apply within 28 days of harvest. Do not graze treated fields.

DANGER TO FISH AND WILDLIFE

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where run-off is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.

RESTRICTIONS

Do not use on crops other than those specified on this label. Use only according to label directions. Applications at rates above those shown may result in illegal crop residues. Do not treat food crops grown in greenhouses.

CONTAINER DISPOSAL

Do not re-use the containers. Rinse equipment and containers and dispose of wastes by burying in non-crop land away from water supplies. Dispose of empty container by burying with wastes or by burning. (Keep out of smoke and fumes).

CAUTION

May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes, and clothing. Do not breathe dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use.

Do not contaminate feed or food. Keep out of reach of children. Not for use or storage in or around the home. If poisoning occurs, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression.

NOTE TO PHYSICIAN: Atropine Sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

CONDITIONS OF SALE

All statements concerning this product or its use apply only when used as directed.

The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

DYLOX is Reg. TM in U. S. and Canadian Pat. Offs. by Farbenfabriken Bayer A.G., Chemagro Corporation licensee.

DRAFT C

71-8412

GEE CHEMICAL CORP. Jacksonville, Florida, U.S.A.