



DIAZINON 14G

GRANULAR INSECTICIDE

ACTIVE INGREDIENT BY WEIGHT
 O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 14.3%
 INERT INGREDIENTS 85.7%

* DIAZINON, registered trademark, CIBA-GEIGY CORPORATION.

CAUTION: MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN.

KEEP OUT OF REACH OF CHILDREN

ACCEPTED
 11/8/74
 UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENT CIDE ACT FOR THE PROTECTION OF POISON REGISTERED UNDER NO. 2342-925 SUBJECT TO ATTACHED COMMENTS

2342-925

Avoid contact with eyes, skin and clothing. Avoid breathing dust. Keep container closed. Use with adequate ventilation. Avoid storage near feed or food products. Wash clothing before reuse. Do not use, pour, spill or store near heat or open flame. Wash thoroughly after handling.

DIRECTIONS FOR USE

Applications to the soil should be worked into the top few inches immediately after application with a rotary hoe, cultivator, disk, narrow or other suitable means to greatest effectiveness.

FIELD AND FORAGE CROP INSECTS

BEANS (Pinto): To control Lesser Cornstalk Borers, apply 7-14 pounds per acre in a 6-inch band centered over the row when beans are 1/2-1 inch high.

CORN: To control Corn Rootworm Larvae, apply 8 1/2-17 ounces per 1,000 linear feet of corn row in a 7-inch band centered over the corn row at planting time or apply amount per acre recommended according to row spacing chart for row treatment.

Corn Rootworms can also be controlled after corn is up. Apply 4 1/2-8 1/2 ounces per 1,000 linear feet of row in a band over the row at first sign of rootworms, but not later than June 14 in the Middle Corn Belt or July 1 in the Northern Corn Belt. Cover treated band with 2-3 inches of soil or apply amount per acre recommended according to row spacing chart for basal treatment. To control European Corn Borers, apply 7-14 pounds per acre over the corn so granules fall into whorls, when 75% of plants show first-generation borers feeding. Make a second application 7 days later, if needed. To control second-generation corn borers, apply 7-14 pounds per acre, as above, when field count shows 100 egg masses per 100 plants. Diazinon 14G n.a. be applied by aircraft for the above use. This material should be uniformly applied to the area to be treated at the recommended rates given above.

Combining Postemergence Treatment for Control of Corn Rootworm Larvae and European Corn Borer North Central United States only: When infestations of these insects coincide, a single applicator may be made to replace the above basal postemergence treatment for Corn Rootworm larvae and one of the above applications for the control of first-generation European Corn Borer. Use the combination treatment where European Corn Borer egg masses or larvae are present at the same time Corn Rootworm Larvae should be controlled. Treatment should be made at the peak of first-generation European Corn Borer egg laying but not later than June 24 in the Middle Corn Belt or July 1 in the Northern Corn Belt for adequate rootworm control.

Apply 7-14 pounds per acre over the corn plants. Cover granules falling to the ground by throwing 2-3 inches of soil into the corn row with a disk tiller or cultivator. Repeat application to corn whorls 7 days later, if needed, for control of European Corn Borer. To control second-generation Corn Borers, apply 7-14 pounds per acre as above, when field count shows 100 egg masses per 100 plants.

To control Fall Armyworms, apply 7-14 pounds per acre over the corn so granules fall into whorls when Armyworms first appear. Repeat application, if necessary.

To control Southwestern Corn Borers, apply 7-14 pounds per acre over the corn so granules fall into whorls when second-generation borers first appear. Repeat application one week later.

NOTE: Do not feed treated fodder to dairy or beef cattle or sheep for 10 days following application. Corn may be picked immediately. Temporary spotting of foliage sometimes occurs following application.

To control Cutworms, broadcast 14-28 pounds per acre. For Surface Cutworms apply just prior to planting or when larvae first appear, and work into soil 1-2 inches. Do not apply after layby. For Subterranean Cutworms, apply just prior to planting and work into soil 3-6 inches.

To control Wireworms, broadcast 21-28 pounds per acre before planting. Work into soil 4-6 inches.

To control Lesser Cornstalk Borers, apply 8 1/2-17 ounces per 1,000 linear feet of row in a 10-inch band centered over the row. Make application at planting time or when crop begins to emerge. Cover the treated band with 1-2 inches of soil or apply amount per acre recommended according to row spacing chart for basal treatment.

To control Seed Corn Maggots, broadcast 14-28 pounds per acre just prior to planting. Work into soil 2 inches.

To control Garden Symphylans (Centipedes), broadcast 70 pounds per acre just prior to planting, and work into soil 4-8 inches. This preplant treatment will also control Cutworms (Surface and Subterranean) and Wireworms.

SUGGESTED RATES ACCORDING TO CORN ROW SPACING

Row spacing	Linear feet of corn row/acre	Amount of Diazinon 14G per acre	
		Row treatment at planting time	Postemergence basal treatment
48"	10,800	6-12 pounds	3-6 pounds
42"	12,455	6 1/2-13 pounds	3 1/2-6 1/2 pounds
40"	13,068	7-14 pounds	3 1/2-7 pounds
38"	13,939	7 1/2-15 pounds	3 1/2-7 1/2 pounds
36"	14,520	8-16 pounds	4-8 pounds
30"	17,424	9 1/2-19 pounds	4 1/2-9 1/2 pounds
24"	21,780	11 1/2-23 pounds	5 1/2-11 1/2 pounds
20"	26,136	14-28 pounds	7-14 pounds
15"	34,848	18 1/2-37 pounds	9 1/2-18 1/2 pounds

Note: For row spacings not listed, calculate the appropriate dosage rates based on the 40-inch row figures.

PEANUTS: To control Southern Corn Rootworms, broadcast 28 pounds per acre or the equivalent in a band over the fruiting zone, just prior to pegging. Work into top few inches of soil. To control Cutworms, apply 14 pounds per acre in a 12-inch band over the row when cutworms first appear. Up to three additional applications at three-week intervals may be necessary. Do not apply within 21 days of peanut harvest. To control Lesser Cornstalk Borers, apply 14-21 pounds per acre in a band over the fruiting zone just prior to pegging. Cover the band with 1-2 inches of soil.

NOTE: Livestock (1) may be fed peanut hulls (2) may be fed peanut forage 7 days following treatment and (3) may be fed peanut hay 21 days following treatment.

SORGHUM: To control Cutworms, broadcast 14-28 pounds per acre just prior to planting. Work into soil 2-3 inches to control Surface Cutworms, 3-6 inches to control Subterranean Cutworms. To control Lesser Cornstalk Borers, apply 7-14 pounds per acre in a 10-inch band over the row at planting time or as crop emerges. Cover the band with 1-2 inches of soil.

NOTE: Do not apply directly to the seed furrow as reduction of sorghum seed germination may occur.

SOYBEANS: To control Cutworms, broadcast 14-28 pounds per acre just prior to planting. Work into soil 2-3 inches to control Surface Cutworms, 3-6 inches to control Subterranean Cutworms. To control Lesser Cornstalk Borers, apply 7-14 pounds per acre in a 6-inch band over the row just prior to planting or when soybeans are 1-2 inches high. Cover the band with 1-2 inches of soil.

TOBACCO: To control Cutworms, broadcast 14-21 pounds per acre before transplanting. Work into soil 2-4 inches.

To control Mole Crickets, apply 7 pounds per acre when insects are present. Make application to plant beds or fields 1-2 days prior to sowing of seed or transplanting, and work into soil 1-2 inches.

To control Seed Corn Maggots, broadcast 10 1/2 pounds per acre 3-4 weeks before transplanting. Work into soil 2 inches.

To control Wireworms, broadcast 14-21 pounds per acre 3-4 weeks before transplanting. Work into soil 6-9 inches.

FORAGE CROPS (Alfalfa, Clover, Cowpeas, and Lespedeza): To control Cutworms, broadcast 14-28 pounds per acre just prior to planting. Work into soil 2-3 inches to control Surface Cutworms, 3-6 inches to control Subterranean Cutworms.

VEGETABLE CROP INSECTS

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, and CAULIFLOWER: To control Root Maggots, broadcast 14-21 pounds per acre before planting or transplanting. Work into soil 3-4 inches.

ONIONS: To control Onion Maggots, apply 14-28 pounds per acre as a broadcast soil treatment just prior to planting. Thoroughly mix the chemical into the top 3-4 inches of soil by disking, harrowing or other suitable means.

RADISHES: To control Root Maggots, apply 10 1/2-14 pounds per acre (2 1/2-2 1/2 ounces per 1,000 linear feet of row) in the seed furrow at planting time.

SWEET CORN: See Corn under Field and Forage Crops (same as field corn).

TOMATOES: To control Vinegar Flies (Drosophila spp.), broadcast 7 pounds per acre when flies first appear. Repeat in 7-10 day intervals or as required.

ADDITIONAL VEGETABLE CROP INSECTS

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CANTALOUPE, CAULIFLOWER, CELERY, COLLARDS, CUCUMBERS, ENDIVE (ESCAROLE), KALE, LETTUCE, LIMA BEANS, MUSKMELONS, PARSLEY, PEAS, PEPPERS, POLE BEANS, SNAP BEANS, SPINACH, SUMMER SQUASH, SWISS CHARD, TOMATOES, WATERMELONS, WINTER SQUASH: To control Cutworms (Surface and Subterranean), broadcast 14-28 pounds per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms, 3-6 inches to control subterranean cutworms.

POTATOES: To control Cutworms (Surface and Subterranean), broadcast 14-28 pounds per acre before planting. Work into soil 4-6 inches.

CABBAGE, CARROTS, LETTUCE, PEAS, POLE BEANS, RADISHES, RED BEETS, SNAP BEANS, TOMATOES, TURNIPS: To control Garden Symphylans (Centipedes), broadcast 70 pounds per acre just prior to planting. Work into soil 4-8 inches. This preplant treatment will also control Cutworms (Surface and Subterranean), Mole Crickets, Root Maggots, and Wireworms.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, COLLARDS, ENDIVE (ESCAROLE), KALE, LETTUCE, MUSTARD, PEPPERS, RADISHES, TOMATOES, TURNIPS: To control Mole Crickets, apply 7 pounds per acre when insects are present. Make application to plant beds or fields 7-21 days prior to sowing of seed or transplanting and work into soil 1-2 inches.

BEANS (Lima, Pole, Snap): To control Lesser Cornstalk Borers, apply 7-14 pounds per acre in a 6-inch band centered over the row. Apply when plants are 1/2-1 inch tall.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CANTALOUPE, CARROTS, CAULIFLOWER, CELERY, COLLARDS, CUCUMBERS, ENDIVE (ESCAROLE), KALE, LETTUCE, LIMA BEANS, MUSKMELONS, ONIONS, PARSLEY, PEAS, PEPPERS, POLE BEANS, SNAP BEANS, SPINACH, SUMMER SQUASH, SWEET POTATOES, SWISS CHARD, TOMATOES, WATERMELONS, WINTER SQUASH: To control Wireworms, broadcast 21-28 pounds per acre just prior to planting. Work into soil 4-8 inches.

POTATOES: To control Wireworms, broadcast 21-42 pounds per acre before planting. Work into soil 2-8 inches or apply 7-21 pounds per acre at planting, placing Diazinon in bands 3-6 inches deep along both sides of row.

SWEET POTATOES (Southeastern U.S.): To control Wireworms, broadcast 21 pounds per acre over the tops of vines when roots begin to enlarge.

SMALL FRUIT CROP INSECTS

STRAWBERRIES: To control Mole Crickets, apply 7 pounds per acre when insects are present. Make application to plant beds or fields 7-21 days prior to transplanting and work into soil 1-2 inches.

LAWN INSECTS

LAWNS: To control Lawn Chinch Bugs, apply 1/2-1 pound per 1,000 square feet, 2 1/2-5 pounds per 5,000 square feet and brush lawn to work granules down to the soil surface. Water lawn thoroughly after application. Repeat application in 7-10 days if necessary. For longer residual control and on dense growing turf, use 1-1 1/2 pounds per 1,000 square feet, 5-7 1/2 pounds per 5,000 square feet.

NOTE: Do not allow children in treated areas until surfaces are dry.

ENVIRONMENTAL PRECAUTIONS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL

Do not reuse container. Destroy it by burying or burning. Avoid smoke or fumes.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. THE MANUFACTURER MAKES NO WARRANTIES, EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE DESCRIPTION ON THE LABEL. Read all directions carefully.

NET WEIGHT 50 POUNDS



DIAZINON 14G

GRANULAR INSECTICIDE

ACTIVE INGREDIENT
O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate
INERT INGREDIENTS

BY WEIGHT
14.3%
85.7%

1 DIAZINON registered trademark. CIBA-GEIGY CORPORATION.

CAUTION: MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN.

KEEP OUT OF REACH OF CHILDREN

ACCEPTED
11/8/74
3342-925

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT REGISTERED UNDER HOWARD COUNTY, MISSOURI TO ASSIGNED COMMENTS

Avoid contact with eyes, skin and clothing. Avoid breathing dust. Keep container closed. Use with adequate ventilation. Avoid storage near feed or food products. Wash clothing before reuse. Do not use, pour, spill or store near heat or open flame. Wash thoroughly after handling.

DIRECTIONS FOR USE

Applications to the soil should be worked into the top few inches immediately after application with a rotary hoe, cultivator, disk, harrow, or other suitable means for greatest effectiveness.

FIELD AND FORAGE CROP INSECTS

BEANS (Pinto) To control Lesser Cornstalk Borers apply 7-14 pounds per acre in a 6-inch band centered over the row when beans are 1/2-1 inch high.

CORN To control Corn Rootworm (larvae) apply 8 1/2-17 ounces per 1,000 linear feet of corn row in a 7-inch band centered over the corn row at planting time or apply amount per acre recommended according to row spacing chart for row treatment. Corn Rootworms can also be controlled after corn is up. Apply 4 1/2-8 1/2 ounces per 1,000 linear feet of row in a band over the row at first sign of rootworms, but not later than June 14 in the Middle Corn Belt or July 1 in the Northern Corn Belt. Cover treated band with 2-3 inches of soil or apply amount per acre recommended according to row spacing chart for basal treatment. To control European Corn Borers apply 7-14 pounds per acre over the corn so granules fall into whorls, when 75% of plants show first general on borers feeding. Make a second application 7 days later, if needed. To control second generation corn borers, apply 7-14 pounds per acre, as above, when field count shows 100 egg masses per 100 plants. Diazinon 14G may be applied by aircraft for the above use. This material should be uniformly applied to the area to be treated at the recommended rates given above.

Combining Postemergence Treatment for Control of Corn Rootworm Larvae and European Corn Borer North Central United States only: When infestations of these insects coincide a single application may be made to replace the above basal postemergence treatment for Corn Rootworm larvae and one of the above applications for the control of first-generation European Corn Borer. Use the combination treatment where European Corn Borer egg masses or larvae are present at the same time Corn Rootworm larvae should be controlled. Treatment should be made at the peak of first-generation European Corn Borer egg laying, but not later than June 24 in the Middle Corn Belt or July 1 in the Northern Corn Belt for adequate rootworm control.

Apply 7-14 pounds per acre over the corn plants. Cover granules falling to the ground by throwing 2-3 inches of soil into the corn row with a disk, harrow or cultivator. Repeat application to corn whorls 7 days later, if needed for control of European Corn Borer. To control second generation Corn Borers, apply 7-14 pounds per acre as above when field count shows 100 egg masses per 100 plants.

To control Fall Armyworms, apply 7-14 pounds per acre over the corn so granules fall into whorls when Armyworms first appear. Repeat application if necessary.

To control Southern Corn Belt Borers, apply 7-14 pounds per acre over the corn so granules fall into whorls when second generation borers first appear. Repeat application one week later.

NOTE: Do not feed treated fodder to dairy or beef cattle or sheep for 10 days following application. Corn may be poked immediately. Temporary spotting of forage sometimes occurs following application.

To control Cutworms, broadcast 14-28 pounds per acre. For Surface Cutworms apply just prior to planting or when larvae first appear, and work into soil 1-2 inches. Do not apply after July 1. For Subterranean Cutworms apply just prior to planting and work into soil 3-6 inches.

To control Wireworms broadcast 21-28 pounds per acre before planting. Work into soil 4-6 inches.

To control Lesser Cornstalk Borers, apply 8 1/2-17 ounces per 1,000 linear feet of row in a 10-inch band centered over the row. Make application at planting time or when crop begins to emerge. Cover the treated band with 1-2 inches of soil or apply amount per acre recommended according to row spacing chart for basal treatment.

To control Seed Corn Maggots, broadcast 14-28 pounds per acre just prior to planting. Work into soil 2 inches.

To control Garden Symphylans (Centipedes), broadcast 70 pounds per acre just prior to planting, and work into soil 4-8 inches. This preplant treatment will also control Cutworms (Surface and Subterranean) and Wireworms.

SUGGESTED RATES ACCORDING TO CORN ROW SPACING			
Row Spacing	Linear feet of corn row per acre	Amount of Diazinon 14G per acre	
		Row treatment at planting time	Postemergence basal treatment
48"	10,890	6 1/2-12 pounds	3-6 pounds
42"	12,455	6 1/2-13 pounds	3 1/2-6 1/2 pounds
36"	13,069	7-14 pounds	3 1/2-7 pounds
30"	15,933	7 1/2-15 pounds	3 1/2-7 1/2 pounds
24"	17,424	8-16 pounds	4-8 pounds
20"	21,780	9-19 pounds	4 1/2-9 1/2 pounds
18"	26,156	11-23 pounds	5 1/2-11 1/2 pounds
15"	34,848	14-28 pounds	7-14 pounds

Note: For row spacings not listed calculate the appropriate dosage rates based on the 40-inch row figures.

PEANUTS To control Southern Corn Rootworms, broadcast 28 pounds per acre or the equivalent in a band over the fruiting zone just prior to pegging. Work into top few inches of soil. To control Cutworms, apply 14 pounds per acre in a 12-inch band over the row when cutworms first appear. Up to three additional applications at three week intervals may be necessary. Do not apply within 21 days of peanut harvest. To control Lesser Cornstalk Borers, apply 14-21 pounds per acre in a band over the fruiting zone just prior to pegging. Cover the band with 1-2 inches of soil. **NOTE:** Livestock (1) may be fed peanut hulls (2) may be fed peanut forage 7 days following treatment and (3) may be fed peanut hay 21 days following treatment.

SORGHUM To control Cutworms, broadcast 14-28 pounds per acre just prior to planting. Work into soil 2-3 inches. To control Surface Cutworms, 3-6 inches to control Subterranean Cutworms, To control Lesser Cornstalk Borers, apply 7-14 pounds per acre in a 10-inch band over the row.

NOTE: Do not feed treated forage to dairy or beef cattle or sheep for 10 days following application.

19 NET WEIGHT 50 POUNDS

MANUFACTURED BY
KERR-MCGEE CHEMICAL CORPORATION
OKLAHOMA CITY, OKLAHOMA

SOYBEANS: To control European Corn Borer, apply 7-14 pounds per acre in a 6-inch band centered over the row when soybeans are 1/2-1 inch high. To control European Corn Borer, apply 7-14 pounds per acre in a 6-inch band centered over the row when soybeans are 1/2-1 inch high. To control European Corn Borer, apply 7-14 pounds per acre in a 6-inch band centered over the row when soybeans are 1/2-1 inch high.

BROCCOLI BRUSSEL SPROUTS: To control European Corn Borer, apply 7-14 pounds per acre in a 6-inch band centered over the row when broccoli brussels are 1/2-1 inch high.

BROCCOLI BRUSSEL SPROUTS: To control European Corn Borer, apply 7-14 pounds per acre in a 6-inch band centered over the row when broccoli brussels are 1/2-1 inch high.

BROCCOLI BRUSSEL SPROUTS: To control European Corn Borer, apply 7-14 pounds per acre in a 6-inch band centered over the row when broccoli brussels are 1/2-1 inch high.

BROCCOLI BRUSSEL SPROUTS: To control European Corn Borer, apply 7-14 pounds per acre in a 6-inch band centered over the row when broccoli brussels are 1/2-1 inch high.

BROCCOLI BRUSSEL SPROUTS: To control European Corn Borer, apply 7-14 pounds per acre in a 6-inch band centered over the row when broccoli brussels are 1/2-1 inch high.

BROCCOLI BRUSSEL SPROUTS: To control European Corn Borer, apply 7-14 pounds per acre in a 6-inch band centered over the row when broccoli brussels are 1/2-1 inch high.



DIAZINON 14G

GRANULAR INSECTICIDE

ACTIVE INGREDIENT
 O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 14.3%
 INERT INGREDIENTS 85.7%

1 DIAZINON, registered trademark, CIBA-GEIGY CORPORATION.

CAUTION: MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN.

KEEP OUT OF REACH OF CHILDREN

ACCEPTED
 11/8/74
 2342-425
 UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR THE PROTECTION OF MAN AND ANIMALS FROM POISON REGISTERED UNDER NO. 2342-425 SUBJECT TO AMENDED COMMENTS.

Avoid contact with eyes, skin and clothing. Avoid breathing dust. Keep container closed. Use with adequate ventilation. Avoid storage near feed or food products. Wash clothing before reuse. Do not use, pour, spill or store near heat or open flame. Wash thoroughly after handling.

19 NET WEIGHT 50 POUNDS

SOYBEANS: To control Cutworms broadcast 14-28 pounds per acre just prior to planting. Work into soil 2-3 inches. To control Surface Cutworms 3-6 inches to control Subterranean Cutworms. To control Lesser Cornstalk Borers apply 7-14 pounds per acre in a 6 inch band over the row just prior to planting or when soybeans are 1-2 inches high. Cover the band with 1-2 inches of soil. ✓
TORACCO: To control Cutworms broadcast 14-21 pounds per acre before transplanting. Work into soil 2-4 inches. To control Mole Crickets, apply 7 pounds per acre when insects are present. Make application to plant beds or fields 1-7 days prior to sowing of seed or transplanting and work into soil 1-2 inches. ✓
 To control Seed Corn Maggots broadcast 10-17 pounds per acre 3-4 weeks before transplanting. Work into soil 2 inches. ✓
 To control Wireworms broadcast 14-21 pounds per acre at planting. Work into soil 6-9 inches. ✓
FORAGE CROPS (Alfalfa, Clover, Grasses, and Lespedeza): To control Cutworms broadcast 14-28 pounds per acre just prior to planting. Work into soil 2-3 inches. To control Surface Cutworms 3-6 inches to control Subterranean Cutworms. ✓

VEGETABLE CROP INSECTS

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, and CAULIFLOWER: To control Root Maggots, broadcast 14-21 pounds per acre before planting or transplanting. Work into soil 3-4 inches. ✓
ONIONS: To control Onion Maggots, apply 14-28 pounds per acre as a broadcast soil treatment just prior to planting. Thoroughly mix the chemical into the top 3-4 inches of soil by disking, harrowing or other suitable means. ✓
RADISHES: To control Root Maggots, apply 10 1/2 - 14 pounds per acre (2 1/2 - 2 3/4 ounces per 1,000 linear feet of row) in the seed furrow at planting time. ✓
SWEET CORN: See Corn under Field and Forage Crops. *(same as field corn)* ✓
TOMATOES: To control Vinegar Flies (Drosophila spp.) broadcast 7 pounds per acre when flies first appear. Repeat in 7-10 day intervals or as required. ✓

ADDITIONAL VEGETABLE CROP INSECTS

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CANTALOUPE, CAULIFLOWER, CELERY, COLLARDS, CUCUMBERS, ENDIVE (ESCAROLE), KALE, LETTUCE, LIMA BEANS, MUSKMELONS, PARSLETS, PEAS, PEPPERS, POLE BEANS, SNAP BEANS, SPINACH, SUMMER SQUASH, SWISS CHARD, TOMATOES, WATERMELONS, WINTER SQUASH: To control Cutworms (Surface and Subterranean), broadcast 14-28 pounds per acre just prior to planting. Work into soil 2-3 inches to control surface cutworms, 3-6 inches to control subterranean cutworms. ✓
POTATOES: To control Cutworms (Surface and Subterranean) broadcast 14-28 pounds per acre just prior to planting. Work into soil 4-6 inches. ✓
CABBAGE, CARROTS, LETTUCE, PEAS, POLE BEANS, RADISHES, RED BEETS, SNAP BEANS, TOMATOES, TURNIPS: To control Garden Symphylans (Centipedes), broadcast 70 pounds per acre just prior to planting. Work into soil 4-8 inches. This pre-plant treatment will also control Cutworms (Surface and Subterranean), Mole Crickets, Root Maggots, and Wireworms. ✓
BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, COLLARDS, ENDIVE (ESCAROLE), KALE, LETTUCE, MUSTARD, PEPPERS, RADISHES, TOMATOES, TURNIPS: To control Mole Crickets, apply 7 pounds per acre when insects are present. Make application to plant beds in fields 1-2 days prior to sowing of seed or transplanting and work into soil 1-2 inches. ✓
BEANS (Lima, Pole, Snap): To control Lesser Cornstalk Borers, apply 7-14 pounds per acre in a 6-inch band centered over the row. Apply when plants are 1/2 - 1 inch tall. ✓
BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CANTALOUPE, CARROTS, CAULIFLOWER, CELERY, COLLARDS, CUCUMBERS, ENDIVE (ESCAROLE), KALE, LETTUCE, LIMA BEANS, MUSKMELONS, ONIONS, PARSLEY, PEAS, PEPPERS, POLE BEANS, SNAP BEANS, SPINACH, SUMMER SQUASH, SWEET POTATOES, SWISS CHARD, TOMATOES, WATERMELONS, WINTER SQUASH: To control Wireworms broadcast 21-42 pounds per acre just prior to planting. Work into soil 4-8 inches. ✓
POTATOES: To control Wireworms, broadcast 21-42 pounds per acre before planting. Work into soil 4-8 inches or apply 7-21 pounds per acre at planting, placing Diazinon in bands 3-6 inches deep along the sides of the row. ✓
SWEET POTATOES (Southeastern U.S.): To control Wireworms, broadcast 21 pounds per acre over the tops of vines when roots begin to enlarge. ✓

SMALL FRUIT CROP INSECTS

STRAWBERRIES: To control Mole Crickets, apply 7 pounds per acre when insects are present. Make application to plant beds or fields 1-2 days prior to transplanting and work into soil 1-2 inches. ✓

LAWN INSECTS

LAWNS: To control Lawn Chinch Bugs, apply 1/2 - 1 pound per 1,000 square feet. 2 1/2 - 5 pounds per 5,000 square feet and brush lawn to work granules down to the soil surface. Water lawn thoroughly after application. Repeat application in 7-10 days if necessary. For longer residual control and on dense growing turf use 1-1 1/2 pounds per 1,000 square feet. 5-7 1/2 pounds per 5,000 square feet. ✓

NOTE: Do not allow children in treated areas until surfaces are dry.

ENVIRONMENTAL PRECAUTIONS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL

Do not reuse container. Destroy it by burning or by other means. Avoid smoke or fumes.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. THE MANUFACTURER MAKES NO WARRANTIES, EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE DESCRIPTION ON THE LABEL. Read all directions carefully.

20

MANUFACTURED BY
 KERR-MCGEE CHEMICAL CORPORATION
 OKLAHOMA CITY, OKLAHOMA