



Product
No. 20-774041

[illegible]

ACTIVE INGREDIENTS:	
Chlorpyrifos Methyl	70.00%
(Chlorpyrifos Methyl) 100% Inert	
INERT INGREDIENTS	30.00%
TOTAL	100.00%

NOTE: The above poundages are suggested on an acre basis, not a gallon basis, and should be applied in enough water to insure thorough distribution over the canopy.

This product mixes well in spray with other commonly used materials such as Dieldrin, Chlorobenzilate, Kelthane, Zineb, Parathion, and Lead Arsenate.

TOMATOES, PEPPERS, POTATOES — Against Early and Late Blight of Potatoes, and Tomatoes, Septoria Leaf Spot of Tomatoes, and Late Blight Spot of Peppers, use 3½ to 4 pounds per 100 gallons of spray every 3 to 7 days as needed.

CUCURBITS (Cucumbers, Watermelons, Cantaloupes, Squash) — Against Downy Mildew use $3\frac{1}{2}$ to 4 pounds per 100 gallons of spray every 5 to 10 days as needed.

CELERY — Against Early Blight use 4 pounds per 100
sq. ft. of spray every 5 to 10 days as needed.

CAUTION: HARMFUL IF SWALLOWED

KEEP OUT OF REACH OF CHILDREN.

CONTAINER DISPOSAL

Do not use as empty container. Destroy by burning
to destroy from smoke and fumes.

COMMITTEE OF ONE

THE USER ASSUMES RESPONSIBILITY FOR THE USE OF THIS PRODUCT AND FOR THE SAFETY OF THE PRODUCT. THE MANUFACTURER MAKES NO WARRANTIES, EXPRESSED OR IMPLIED, WHICH EXTEND BEYOND THE INFORMATION ON THE LABEL. THE MANUFACTURER SHALL IN NO EVENT BE LIABLE FOR CONSEQUENTIAL DAMAGES. READ AND FOLLOW DIRECTIONS CAREFULLY.

EPA R. No.

1-73 W

EPH Est. 2342-FL-1

Monitored by

TEA-MODE CHEMICAL CORP.

Seville, Florida, U.S.A.