



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

November 12, 2025

Kara Welch
Regulatory Affairs Manager
Chr. Hansen, Inc.
9015 West Maple Street
Milwaukee, WI 53214-4298

Subject: Pesticide Registration Improvement Act (PRIA) Labeling Amendment – Revisions to
Update the Expiration Date, Change the File Symbol to an EPA Registration Number,
and Add Referral Statements
Response to the Storage Stability and Corrosion Characteristics Term of Registration
Product Name: Kansas 3 WP
EPA Registration Number: 2375-7
EPA Receipt Date: 7/30/2025
Action Case Number: 00663336

Dear Kara Welch:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable.

Besides the storage stability and corrosion characteristics data term (see below), this approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 § CFR 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S. Environmental Protection Agency. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should EPA find or if it is brought to our attention that a website contains statements or claims substantially

differing from statements or claims made in connection with obtaining a FIFRA section 3 registration, the website will be referred to EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

Additionally, EPA rated submitted storage stability and corrosion characteristics data (MRID No. 526433-01) as acceptable, and Kansas 3 WP is stable up to 24 months. There were no observable differences in the physical changes of the test substance or concentration of the active ingredients. Thus, you have fulfilled term #3 of the September 27, 2024, Kansas 3 WP Registration Notice.

If you have any questions, please contact Amy Krause via email at krause.amy@epa.gov.

Sincerely,



Digitally signed by
JEANNINE KAUSCH
Date: 2025.11.12
13:53:49 -05'00'

Jeannine Kausch, Senior Advisor
Immediate Office
Biopesticides and Pollution
Prevention Division (7511M)
Office of Pesticide Programs

KANSAS 3 WP

Wettable Powder MASTER LABEL

Sub-Label A: Seed Treatment Use

Sub-Label B: Soil-Applied Use

ACCEPTED

Nov 12, 2025

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 2375-7

A wettable powder biological fungicide/nematicide for seed treatment and soil-applied use for early season protection against labeled fungal diseases and soil nematodes.

EPA Reg. No. 2375-7

EPA Est. No. 2375-xx-xxxxx

Active Ingredients:	By Wt.
<i>Bacillus subtilis</i> strain CH4000*	20.0%
<i>Bacillus paralicheniformis</i> strain CH0273*	20.0%
<i>Bacillus paralicheniformis</i> strain CH2970*	20.0%
Other Ingredients:	40.0%
Total	100.0%

*Contains a minimum of 1.0×10^{11} colony forming units (CFU) of each active ingredient per g of product.

KEEP OUT OF REACH OF CHILDREN

[See other panels for additional precautionary information.]

[See complete Precautionary Statements and Directions for Use in booklet inside box on the bag or pouch]

[See complete Precautionary Statements and Directions for Use in booklet below]

Not for sale or use after [Date stamped/placed on labeling will be 24 months after the date of manufacture]

Net Weight: xxxxx

Sold By:
Chr. Hansen
9015 W Maple St.
Milwaukee, WI 53214

Batch/Lot code: _____
[For nonrefillable containers only. Will be located on
this label or on the physical container.]

KANSAS 3 WP

Wettable Powder LABEL

Sub-Label A

Seed Treatment Use

A WP biological fungicide/nematicide for seed treatment use for early season protection against labeled fungal diseases and soil nematodes.

EPA Reg. No. 2375-7

EPA Est. No. 2375-xx-xxxxx

Active Ingredients:	By Wt.
<i>Bacillus subtilis</i> strain CH4000*	20.0%
<i>Bacillus paralicheniformis</i> strain CH0273*	20.0%
<i>Bacillus paralicheniformis</i> strain CH2970*	20.0%
Other Ingredients:	40.0%
Total	100.0%

*Contains a minimum of 1.0×10^{11} colony forming units (CFU) of each active ingredient per g of product.

KEEP OUT OF REACH OF CHILDREN

[See side/back panel for First Aid]

[See complete Precautionary Statements and Directions for Use in booklet inside box on the bag or pouch]

[See complete Precautionary Statements and Directions for Use in booklet below]

Not for sale or use after [Date stamped/placed on labeling will be 24 months after the date of manufacture]

Net Weight: xxxxxx

Sold By:
Chr. Hansen
9015 W Maple St.
Milwaukee, WI 53214

Batch/Lot code: _____
[For nonrefillable containers only. Will be located on
this label or on the physical container.]

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information on this pesticide product (including general health concerns or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday, 8:00AM to 12:00PM Pacific Standard Time (NPIC Website: www.npic.orst.edu). In the event of a medical emergency, call your local poison control center at 1-800-222-1222.</p>	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- protective eyewear
- long-sleeved shirt
- long pants
- shoes plus socks

Mixers/loaders and applicators must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R, or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R, or P filter; OR a NIOSH-approved powered air-purifying respirator with an HE filter. Repeated exposures to high concentrations of microbial proteins can cause allergic sensitization.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d) and (e)], the handler PPE requirements may be reduced or modified as specified in the WPS

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for “applicators and other handlers” and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

User Safety Recommendations

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove clothing/PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

For container sizes <5 gallons

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

For container sizes ≥5 gallons

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval, and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water) is:

- Protective eyewear
- Coveralls
- Chemical-resistant gloves (made of any waterproof material)
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treatment area until seeds have dried or been packaged.

Product Information

KANSAS 3 WP is a biofungicide and bionematicide seed treatment wettable powder formulation that provides suppression and control of labeled nematodes and fungi in a variety of horticultural and broadacre crops for any and all seed practices including farm, greenhouse, nursery, or any other seed development and application processes. Seed must be sound and well-cured before treatment.

Seed Treatment Equipment

KANSAS 3 WP may be used both for commercial and for on-farm application. KANSAS 3 WP can be applied with mechanical, slurry, or mist-type seed treating equipment, as long as the equipment can be calibrated to accurately and uniformly apply the product to seed without undue mechanical damage to the seed. Uniform application to seed is important for all seed treatment products.

Application Instructions

Seed must be sound and well-cured before treatment. Refer to COMPATIBILITY AND TANK MIXING INSTRUCTIONS below and follow the crop-specific the application rates below.

COMPATIBILITY AND TANK MIXING INSTRUCTIONS

COMPATIBILITY

KANSAS 3 WP is physically compatible with many commonly used fungicides, herbicides, insecticides, and(or) nutrient products. However, the compatibility of KANSAS 3 WP with all potential tank-mix partners has not been fully investigated. If tank mixing with other products is desirable, conduct a jar test with the volumes and rates typically used in labeled applications. Using a small container of water, add the proportionate amounts of the products in the following order: wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 15 minutes. Look for signs of separation, globules, sludge, flakes, or other precipitates. Physical compatibility is indicated if the combination remains mixed or can be remixed readily.

TANK MIXING INSTRUCTIONS

When tank-mixing KANSAS 3 WP with other registered pesticides and(or) other products, always read and follow all use directions, restrictions, and precautions of both KANSAS 3 WP and the tank-mix partner(s). It is the pesticide user's responsibility to ensure that all tank-mix products are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Use of the resulting tank mix must be in accordance with the more restrictive label limitations and precautions. DO NOT exceed label dosage rates.

All applications of KANSAS 3 WP involve the following general tank mixing procedures. Begin with clean supply, mixing and application equipment. Add one-half of the required amount of non-chlorinated water to the supply/mixing tank and start agitation. Dissolve the needed amount of KANSAS 3 WP into a small volume of water to make a pre-mix slurry. Add needed amounts of products to the mix tank in the following order: water conditioners, water soluble packages (wait for them to completely dissolve), wettable powders and water-dispersible granular products, followed by liquids, including, liquid slurries (add KANSAS 3 WP pre-mix slurry here), flowables, suspension concentrates, emulsifiable concentrates, with surfactants and(or) adjuvants added last. After adding the required quantity of KANSAS 3 WP and other products to the mix tank, complete filling with water to the required final volume and maintain agitation throughout application. DO NOT allow tank mixture to remain in the tank overnight. DO NOT allow tank mixture to sit for long

Note to Reviewer: Language within brackets is optional label language. Kansas 3 WP Reg. No. 2375-7 Master Label – dated June 27, 2025

periods during the day without agitation. If non-chlorinated water is not available, do not let KANSAS 3 WP remain in chlorinated water for long periods of time as it may have detrimental effect on the bacterial endospores. Follow the advice of your State Cooperative Extension Service for tank mixing procedures with other products.

The appropriate volume of slurry for seed treatment depends on crop, temperature, humidity, type of equipment, and other factors and must be adjusted per normal treating practices for the circumstances. Contact your local supplier or distributor representative for additional specifications for slurry volumes and questions on any tank mixes or restrictions.

Recalibrate equipment to compensate for the required slurry rate to ensure all products are applied at the correct application rate.

NOTE: This product does not contain dye and is not covered by an appropriate tolerance, tolerance exemption, or other clearance under the Federal Food, Drug and Cosmetic Act. To comply with 40 CFR 153.155, therefore, all seed treated commercially with this product must be colored with an EPA-approved dye or colorant of a suitable color to prevent accidental use as food for man or feed for animals.

SEED BAG LABEL REQUIREMENTS

The Federal Seed Act requires that bags containing seeds treated with this product shall be labeled with the following information:

- This seed has been treated with *Bacillus subtilis* strain CH4000 and *Bacillus paralicheniformis* strains CH0273 and *Bacillus paralicheniformis* strains CH2970.
- Do not use for feed, food or oil purposes. Store away from feed and food stuffs.

User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

Do not use KANSAS 3 WP in combination with other seed treatment products unless compatibility has been verified. Read and carefully follow all label directions of each product used in the treating mixture. When using combinations of products, the most restrictive of label limitations and precautions must be followed.

Application Rates, Crops, and Pests

Application Rates

KANSAS 3 WP may be applied as a seed treatment at a product rate of 0.8 to 8.0 oz/100 pounds of seed, following the specific crop directions. Use the lower application rate for light infestations and the higher labeled rate for heavy infestations

Specific Crop Directions

Crop Rotation Restrictions

There are no crop rotation restrictions. Any crop can be rotated at any time after application of KANSAS 3 WP

Crop Group 18 - Nongrass Animal Feeds (Forage, Fodder, Straw, and Hay) Group: Alfalfa, Clover, and Forages		
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*] <i>Globodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	0.5 – 3.0 g/kg or 0.8 – 4.8 oz/100 lbs

*Not registered for use in California

Crop Group 1 - Root and Tuber Vegetables Group: Artichoke		
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.0 – 5.0 g/kg or 1.6 – 8.0 oz/100 lbs

*Not registered for use in California

Crop Group 5 - <i>Brassica</i> (Cole) Leafy Vegetables Group: Broccoli, Cabbage, Cauliflower, Brussels Sprouts, Collards, Kale, Mustard Greens, Kohlrabi, and other <i>brassica</i> crops		
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.0 – 5.0 g/kg or 1.6 – 8.0 oz/100 lbs

*Not registered for use in California

Crop Group 15 -	Cereal Grains Group: (Exclude Corn) Barley, Buckwheat, Millet, Oat, Pearl Millet, Proso Millet, Rice**, Rye, Sorghum, Teosinte, Triticale, Wheat, and other cereal grain Crops	
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.0 – 5.0 g/kg or 1.6 – 8.0 oz/100 lbs

*Not registered for use in California

**Do not apply to flooded fields

Crop Group 15 -	Cereal Grains Group: Corn (Field corn, Popcorn, Sweet corn, and Corn Grown for Seed)	
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp. [*]	1.0 – 5.0 g/kg or 1.6 – 8.0 oz/100 lbs

*Not registered for use in California

Crop Group 20 -	Oilseed Group: Cottonseed	
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Rotylenchulus reniformis</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.0 – 5.0 g/kg or 1.6 – 8.0 oz/100 lbs

*Not registered for use in California

Crop Group 9 -	Cucurbit Vegetables Group: Chayote (fruit), Chinese wax gourd (Chinese preserving melon), Citron melon, Cucumber, Cantaloupe, Edible Gourd (such as Chinese Okra, Cucuzza, Hechima, and Hyotan), Gherkin, Melon, <i>Momordica</i> spp. (such as Balsam apple, Balsam pear, Bitter melon, and Cucumber), Muskmelon, Pumpkin, Summer and Winter Squash (such as Acorn, Butternut, Calabaza, Crookneck, Hubbard, Scallop, Spaghetti, and Zucchini), Squash, Watermelon, Chinese Watermelon, and other cucurbit crops	
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.0 – 5.0 g/kg or 1.6 – 8.0 oz/100 lbs

*Not registered for use in California

Crop Group 8 -	Fruiting Vegetables Group: African eggplant, Bell pepper, Bush tomato, Cocona, Currant tomato, Eggplant, Garden huckleberry, Goji berry, Groundberry, Martynia, Naranjilla, Nonbell pepper, Okra, Pea eggplant, Pepino, Roselle, Scarlet eggplant, Sunberry, Tomatillo, and Tomato	
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*] <i>Globodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.0 – 5.0 g/kg or 1.6 – 8.0 oz/100 lbs

*Not registered for use in California

Crop Group 4 -	Leafy Vegetables (Except <i>Brassica</i> Vegetables) Group: Lettuce, Celery, Spinach, Parsley, Radicchio, and other leafy vegetables crops including those grown for seed production	
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.0 – 5.0 g/kg or 1.6 – 8.0 oz/100 lbs

*Not registered for use in California

Crop Group 6 -	Legume Vegetables (Succulent or Dried): (Exclude Lentils and Soybeans) Beans, Snap beans, Garbanzo beans, Lima beans, Peas, Chick peas, Dried bean and pea (<i>Lupinus</i> spp., <i>Phaseolus</i> spp., <i>Vigna</i> spp., and <i>Pisum</i> spp.), and other legume vegetable crops including those grown for seed production	
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	0.5 – 3.0 g/kg or 0.8 - 4.8 oz/100 lbs

*Not registered for use in California

Crop Group 6 -	Legume Vegetables (Succulent or Dried): Lentils	
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	0.5 – 3.0 g/kg or 0.8 - 4.8 oz/100 lbs

*Not registered for use in California

Crop Group 20 -	Oilseed Group: (Excludes Cottonseed, Peanut, and Soybeans) Canola, Castor, Flax, Palm, Olive, Rapeseed, Safflower, Sesame, Sunflower, and other oil seed crops	
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.0 – 5.0 g/kg or 1.6 – 8.0 oz/100 lbs

*Not registered for use in California

Crop -	Peanut	
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Belonolaimus</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	0.5 – 3.0 g/kg or 0.8 - 4.8 oz/100 lbs

*Not registered for use in California

Note to Reviewer: Language within brackets is optional label language. Kansas 3 WP Reg. No. 2375-7 Master Label – dated June 27, 2025

Crop Group 1 -	Root and Tuber Vegetables Group: Potato	
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	0.5 – 3.0 g/kg or 0.8 - 4.8 oz/100 lbs

*Not registered for use in California

Crop Group 1 -	Root and Tuber Vegetables Group: (Exclude Potato, Sugar Beet, and Artichoke) Arracacha, Arrowroot, Bean and True Yam, Beets, Bitter and Sweet Cassava, Black and Spanish Salsify, Carrot, Celeriac (Celery Root), Chayote (Root), Chicory, Chufa, Dasheen (Taro), Edible Burdock, Edible Cana, Ginger, Ginseng, Green Beet, Horseradish, Leren, Oriental Radish, Parsnip, Radish, Rutabaga, Salsify (Oyster plant), Skirret, Sweet Potato, Tanier (Cocoyam), Turmeric, Turnip, Turnip-rooted Chervil, and Turnip-rooted Parsley, and other root and tuber vegetable crops	
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.0 – 5.0 g/kg or 1.6 – 8.0 oz/100 lbs

*Not registered for use in California

Crop Group 6 -	Legume Vegetables (Succulent or Dried): Soybeans	
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	0.5 – 3.0 g/kg or 0.8 - 4.8 oz/100 lbs

*Not registered for use in California

Crop Group 1 -	Root and Tuber Vegetables Group: Sugarbeet	
Target Nematode	Target Plant Pathogens	Application Rate: Product/Seed Wt:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	0.5 – 3.0 g/kg or 0.8 - 4.8 oz/100 lbs

*Not registered for use in California

USE RESTRICTIONS

Do not use treated seed for food, feed or oil purposes. For at-planting seed treatments, treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Care must be exercised in the handling of treated seed. Augers used for handling treated seed must not be used to move seed for feed, food or oil processing. Do not re-use bags from treated seed to handle food or feed products.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store in a cool, dry place under low humidity. Avoid direct sunlight and avoid excess heat. Keep out of reach of children and animals. Store in original container only. Carefully open container. After partial use, replace lid of container and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal. In case of spill, avoid contact, isolate area, and keep out animals and unprotected persons. Confine spills. Call CHEMTREC (Transportation and Spills) at 1-800-424-9300. To confine spill, dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. If a leaky container must be contained within another, mark the outer container to identify the contents.

Pesticide Disposal

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling

Nonrefillable Plastic Containers (Containers equal to or less than 5 gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank, and drain for 10 seconds after the flow begins to drip. Add water – at least 2% of the container volume, and up to $\frac{1}{3}$ of the volume of water needed to make the proper slurry composition with a maximum of $\frac{1}{4}$ of the container volume, and recap. Shake for 30 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water. Then offer container for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill or by incineration. If burned, stay out of smoke.

Nonrefillable Plastic Containers (Containers greater than 5 gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Add water – at least 2% of the container volume, and up to $\frac{1}{3}$ of the volume of water needed to make the proper slurry composition with a maximum of $\frac{1}{4}$ of the container volume. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 60 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water. Then offer container for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill or by incineration. If burned, stay out of smoke.

Conditions of Sale and Limitation of Warranty and Liability

Notice: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of Chr. Hansen or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and, to the extent consistent with applicable law, Buyer and User agree to hold Chr. Hansen and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the compositional description on the label and is reasonably fit for the purposes stated in the Directions for Use when used in accordance with the directions under normal conditions of use.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CHR. HANSEN MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Chr. Hansen, and, to the extent consistent with applicable law, Buyer assumes the risk of any such use.

Treatment of seed, especially seed that is mechanically damaged or seed known to be of low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small test sample of seed before treating commercial quantities with a selected treatment that includes this product. Due to seed quality and seed storage conditions beyond the control of Chr. Hansen, Chr. Hansen makes no claims or guarantees as to germination of carry-over seed.

To the extent consistent with applicable law, Chr. Hansen or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

Chr. Hansen – Trademark of Chr. Hansen Corporation
© 2020 Chr. Hansen Corporation All rights reserved.

KANSAS 3 WP

Wettable Powder LABEL

Sub-Label B

Soil-applied Use

A wettable powder biological fungicide/nematicide for soil-applied use for protection against listed fungal diseases and soil nematodes.

EPA Reg. No. 2375-7

EPA Est. No. 2375-xx-xxxxx

Active Ingredients:	By Wt.
<i>Bacillus subtilis</i> strain CH4000*	20.0%
<i>Bacillus paralicheniformis</i> strain CH0273*	20.0%
<i>Bacillus paralicheniformis</i> strain CH2970*	20.0%
Other Ingredients:	40.0%
Total	100.0%

*Contains a minimum of 1.0×10^{11} colony forming units (CFU) of each active ingredient per g of product.

KEEP OUT OF REACH OF CHILDREN

[See side/back panel for First Aid]

[See complete Precautionary Statements and Directions for Use in booklet inside box on the bag or pouch]

[See complete Precautionary Statements and Directions for Use in booklet below]

Not for sale or use after [Date stamped/placed on labeling will be 24 months after the date of manufacture]

Net Weight: XXXX

Sold By:
Chr. Hansen
9015 W Maple St.
Milwaukee, WI 53214

Batch/Lot code: _____
[For nonrefillable containers only. Will be located on
this label or on the physical container.]

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information on this pesticide product (including general health concerns or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday 8:00AM to 12:00PM Pacific Standard Time (NPIC Website: www.npic.orst.edu). In the event of a medical emergency, call your local poison control center 1-800-222-1222.	

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- protective eyewear
- long-sleeved shirt
- long pants
- shoes plus socks

Mixers/loaders and applicators must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R, or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R, or P filter; OR a NIOSH-approved powered air-purifying respirator with an HE filter. Repeated exposures to high concentrations of microbial proteins can cause allergic sensitization.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d) and (e)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

User Safety Recommendations

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove clothing/PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

For container sizes <5 gallons

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

For container sizes ≥5 gallons

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water) is:

- Protective eyewear
- Coveralls
- Chemical resistant gloves (made of any waterproof material)
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treatment area until applied product is covered or dried in soil.

PRODUCT INFORMATION

KANSAS 3 WP is a Biofungicide/Bionematicide soil applied wettable powder formulation for the suppression and control of labeled fungi and soil nematodes in a variety of horticultural and broadacre crops.

SOIL APPLICATION

KANSAS 3 WP Biofungicide/Bionematicide can be used on labeled crops through direct soil application by drench, side-dress, in-furrow, or by chemigation through irrigation. The rate of application is variable according to pest pressure. Use lower labeled rates under light to moderate pest infestations, and higher labeled rates under heavier pest pressure. Use the highest labeled rate under arid conditions.

DRENCH, SIDE-DRESS, OR IN-FURROW APPLICATION INSTRUCTIONS

After conducting initial tank mixing procedures (see COMPATIBILITY AND TANK MIXING section), apply KANSAS 3 WP Biofungicide/Bionematicide directly to soil by drench, side-dress or in-furrow (either T-band or directly over seed in the open furrow). This product may be applied as a soil treatment in greenhouses. After such applications, ensure the product is covered with soil.

CHEMIGATION THROUGH IRRIGATION INSTRUCTIONS

After conducting initial tank mixing procedures (see COMPATIBILITY AND TANK MIXING section) apply KANSAS 3 WP Biofungicide/Bionematicide only through center pivot, motorized-lateral move, traveling gun, solid set or portable (wheel move, side roll, end tow, or hand move) and drip irrigation systems, **as outlined in Sections A and B below**. Chemigation systems must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. Determine which type of irrigation system is in place, then refer to the appropriate directions provided below for each type (Sections A and B below). See crops section on the label for required treatment rates and additional use information. **DO NOT** apply KANSAS 3 WP Biofungicide/Bionematicide through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated, water-based tank mixes. If you have questions about calibration of irrigation equipment, contact State Extension Service specialists, equipment manufacturers or other experts. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, must shut the system down and make necessary adjustments should the need arise. KANSAS 3 WP Biofungicide/Bionematicide has not been sufficiently tested when applied through irrigation systems to assure consistent product performance for all labeled uses. The following application techniques are provided for user reference but **DO NOT** constitute a warranty of fitness for application through irrigation equipment.

DO NOT connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label prescribed safety devices for public water systems are in place. 'Public water system' means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), back flow preventer, or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an alternative to the RPZ, the water from the public water system must be discharged into a reservoir tank prior to pesticide introduction. There must be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid

Note to Reviewer: Language within brackets is optional label language. Kansas 3 WP Reg. No. 2375-7 Master Label – dated June 27, 2025

back toward the injection pump. Pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The systems must contain functional interlocking controls, to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. DO NOT apply when wind speed favors drift beyond the area intended for treatment. Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur.

KANSAS 3 WP Biofungicide/Bionematicide can be applied to labeled crops as a liquid tank-mix to the soil by drench, side dress, in-furrow (as a T-band or over the open furrow), or by chemigation through irrigation systems. The rate of application is variable according to pest pressure. Use lower labeled rates under light to moderate pest infestations, and higher labeled rates under heavier pest pressure. Use the highest labeled rate under arid conditions.

SECTION A. INSTRUCTIONS FOR CENTER PIVOT, MOTORIZED-LATERAL MOVE AND TRAVELING GUN IRRIGATION EQUIPMENT

After tank-mixing procedures, KANSAS 3 WP Biofungicide/Bionematicide can be applied through center pivot, motorized-lateral move, or traveling gun systems. These continuously moving systems must use a positive displacement injection pump of either diaphragm or piston type and be constructed of materials that are compatible with pesticides. They must also be capable of being fitted with a system interlock and capable of injection at pressures approximately 2-3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems. Thoroughly mix required amount of KANSAS 3 WP Biofungicide/Bionematicide for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until KANSAS 3 WP Biofungicide/Bionematicide has been cleared from the last sprinkler head.

SECTION B. INSTRUCTIONS FOR SOLID-SET, PORTABLE (WHEEL MOVE, SIDE ROLL, END TOW, OR HAND MOVE) AND DRIP IRRIGATION SYSTEMS

For Solid-Set, Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Systems

After tank-mixing procedures, KANSAS 3 WP Biofungicide/Bionematicide can be applied through stationary irrigation systems, including Solid-Set, Portable (Wheel Move, Side Roll, End Tow, or Hand Move). With these systems, an effectively designed in-line Venturi applicator unit is preferred to support even and quick distribution of the product. However, a positive-displacement pump can also be used. For solid set systems, determine acreage covered by sprinklers. Fill the tank of injection equipment with water and adjust flow to use contents over 30 to 45 minutes. Mix desired amount of this product for acreage to be covered with water so that the total mixture of this product plus water in the injection tank is equal to the quantity of water used during calibration. Provide chemical supply tank agitation sufficient for mixing until chemigation is completed. Operate entire system at normal pressures as advised by the manufacturer of injection equipment used, for amount of time established during calibration. This product can be injected during the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and

continue to operate irrigation system until the product(s) has been cleared from the lines and last sprinkler head.

For drip irrigation systems

After tank-mixing procedures, KANSAS 3 WP Biofungicide/Bionematicide can be applied through stationary drip irrigation systems. Provide chemical supply tank agitation sufficient for mixing until chemigation is completed. With these systems, an effectively designed in-line Venturi applicator unit is preferred to support even and quick distribution of the product. Fill the tank of injection equipment with water and adjust the flow out of the supply/mix tank after 25% of irrigation water allocation has been applied in order to ensure the irrigation lines are primed and a wetting front in the roots has been created. Then, apply the KANSAS 3 WP Biofungicide/Bionematicide mixture into 50% of the irrigation water allocation volume to ensure optimal distribution of the product uniformly in the crop. Then, follow up with 25% of the total irrigation water allocation volume to ensure the product(s) have been cleared from irrigation lines. Operate entire system at normal pressures as advised by the manufacturer of injection equipment used for the amount of time established during calibration.

COMPATIBILITY AND TANK MIXING INSTRUCTIONS

COMPATIBILITY

KANSAS 3 WP Biofungicide/Bionematicide is physically compatible with many commonly used fungicides, herbicides, insecticides, and(or) nutrient products. However, the compatibility of KANSAS 3 WP Biofungicide/Bionematicide with all potential tank-mix partners has not been fully investigated. If tank mixing with other products is desirable, conduct a jar test with the volumes and rates typically used in labeled applications. Using a small container of water, add the proportionate amounts of the products in the following order: wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 15 minutes. Look for signs of separation, globules, sludge, flakes, or other precipitates. Physical compatibility is indicated if the combination remains mixed or can be remixed readily.

TANK MIXING INSTRUCTIONS

When tank-mixing KANSAS 3 WP Biofungicide/Bionematicide with other registered pesticides and(or) other products, always read and follow all use directions, restrictions, and precautions of both KANSAS 3 WP Biofungicide/Bionematicide and the tank-mix partner(s). It is the pesticide user's responsibility to ensure that all tank-mix products are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Use of the resulting tank mix must be in accordance with the more restrictive label limitations and precautions. DO NOT exceed label dosage rates.

All applications of KANSAS 3 WP Biofungicide/Bionematicide involve the following general tank mixing procedures. Begin with clean supply, mixing and application equipment. Add one-half of the required amount of non-chlorinated water to the supply/mixing tank and start agitation. Dissolve the needed amount of KANSAS 3 WP Biofungicide/Bionematicide into a small volume of water to make a pre-mix slurry. Add needed amounts of products to the mix tank in the following order: water conditioners, water soluble packages (wait for them to completely dissolve), wettable powders and water-dispersible granular products, followed by liquids, including, liquid slurries (add KANSAS 3 WP Biofungicide/Bionematicide pre-mix slurry here), flowables, suspension concentrates, emulsifiable concentrates, with surfactants and(or) adjuvants added last. After adding the required quantity of KANSAS 3 WP Biofungicide/Bionematicide and other products to the mix tank, complete filling with water to the required final volume and maintain agitation

throughout application. DO NOT allow tank mixture to remain in the tank overnight. DO NOT allow tank mixture to sit for long periods during the day without agitation. If non-chlorinated water is not available, DO NOT let KANSAS 3 WP Biofungicide/Bionematicide remain in chlorinated water for long periods of time as it may have detrimental effect on the bacterial endospores. Do not combine KANSAS 3 WP Biofungicide/Bionematicide with pesticides, surfactants, or fertilizers with which there has been no previous experience or use demonstrating that they are physically compatible, effective, and non-injurious under your use conditions. Follow the advice of your State Cooperative Extension Service for tank mixing procedures with other products.

Application Rates, Crops, and Pests

Specific Crop Directions

Crop Rotation Restrictions

There are no crop rotation restrictions. Any crop can be rotated at any time after application of KANSAS 3 WP

Application Rates

KANSAS 3 WP may be applied as a soil applied fungicide/nematicide treatment at rates between 1.43 - 5.71 oz/A in 3 – 22 gal total solution for the following crops and fungi/nematode pests

Crop Group 18 -		Nongrass Animal Feeds (Forage, Fodder, Straw, and Hay) Group: Alfalfa, Clover, and Forages	
Target Nematode	Target Plant Pathogens	Application Rate:	
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*] <i>Globodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A	

*Not registered for use in California

Crop Group 11 -		Pome Fruits Group: Apple, Crabapple, Pear, Quince, Mayhaw, and other pome fruit	
Target Nematode	Target Plant Pathogens	Application Rate:	
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*] <i>Xiphinema</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A	

*Not registered for use in California

Crop Group 12 -	Stone Fruit Group: Apricot, Cherry, Nectarine, Peach, Plum, and other stone fruit crops	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 1 -	Root and Tuber Vegetables Group: Artichoke	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 22 -	Stalk, Stem and Leaf Petiole Vegetable Group: Asparagus	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 24 -	Tropical and Subtropical Fruit, Inedible Peel Group: Avocado and Mango	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 5 -	Brassica (Cole) Leafy Vegetables Group: Broccoli, Cabbage, Cauliflower, Brussels Sprouts, Collards, Kale, Mustard Greens, Kohlrabi, and other <i>brassica</i> crops	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 24 -	Tropical and Subtropical Fruit, Inedible Peel Group: Banana, Plantain, and Pineapple	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 15 -	Cereal Grains Group: (Exclude Corn) Barley, Buckwheat, Millet, Oat, Pearl Millet, Proso Millet, Rice**, Rye, Sorghum, Teosinte, Triticale, Wheat, and other cereal grain crops	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

**Do not apply to flooded fields

Crop Group 15 -	Cereal Grains Group: Corn (Field corn, Popcorn, Sweet corn, and Corn Grown for Seed)	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 20 -	Oilseed Group: Cottonseed	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Rotylenchulus reniformis</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 10 -	Citrus Fruit Group: Calamondin, Citrus citron, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Sour and Sweet Orange, Pummelo, Satsuma mandarin, and other citrus fruit crop	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop -	Coffee	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 9 -	Cucurbit Vegetables Group: Chayote (fruit), Chinese wax gourd (Chinese preserving melon), Citron melon, Cucumber, Cantaloupe, Edible Gourd (such as Chinese Okra, Cucuzza, Hechima, and Hyotan), Gherkin, Melon, <i>Momordica</i> spp. (such as Balsam apple, Balsam pear, Bitter melon, and Cucumber), Muskmelon, Pumpkin, Summer and Winter Squash (such as Acorn, Butternut, Calabaza, Crookneck, Hubbard, Scallop, Spaghetti, and Zucchini), Squash, Watermelon, Chinese Watermelon, and other cucurbit crops	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 8 -	Fruiting Vegetables Group: African eggplant, Bell pepper, Bush tomato, Cocona, Currant tomato, Eggplant, Garden huckleberry, Goji berry, Groundberry, Martynia, Naranjilla, Nonbell pepper, Okra, Pea eggplant, Pepino, Roselle, Scarlet eggplant, Sunberry, Tomatillo, and Tomato	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*] <i>Globodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 13-07 -	Berry and Small Fruit Crop Group: Grapes	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*] <i>Tylenchulus semipenetrans</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop -	Hemp	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 19 -	Herbs and Spices Group: Fresh and Dried Basil, Black Pepper, Chive, Celery seed, Dill seed, and other herbs and spices	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop -	Hops	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 13-07 -	Berry and Small Fruit Crop Group: Kiwi	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 4 -	Leafy Vegetables (Except <i>Brassica</i> Vegetables) Group: Lettuce, Celery, Spinach, Parsley, Radicchio, and other leafy vegetables crops including those grown for seed production	
Target Nematode	Target Plant Pathogens	Application Rate
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 6 -	Legume Vegetables (Succulent or Dried): (Exclude Lentils and Soybeans) Beans, Snap beans, Garbanzo beans, Lima beans, Peas, Chick peas, Dried bean and pea (<i>Lupinus</i> spp., <i>Phaseolus</i> spp., <i>Vigna</i> spp., and <i>Pisum</i> spp.), and other legume vegetable crops including those grown for seed production	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 6 -	Legume Vegetables (Succulent or Dried): Lentils	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop 25 -	Mint	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 20 -	Oilseed Group: (Excludes Cottonseed, Peanut, and Soybeans) Canola, Castor, Flax, Palm, Olive, Rapeseed, Safflower, Sesame, Sunflower, and other oil seed crops	
Target Nematode	Target Plant Pathogens	Application Rate
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 3 -	Bulb Vegetables (<i>Allium</i> SPP.) Group: Onion, Garlic, Shallots, and other bulb vegetables including those grown for seed production	
Target Nematode	Target Plant Pathogens	Application Rate
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop -	Ornamental Flowers	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop -	Peanut	
Target Nematode	Target Plant Pathogens	Application Rate:
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Belonolaimus</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 23 -	Tropical and Subtropical Fruit, Edible Peel Group: Olive and Palm Tree	
Target Nematode	Target Plant Pathogens	Application Rate
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*] <i>Hemicriconemoides</i> [*] <i>Rotylenchulus</i> [*] <i>Tylenchulus</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 24 -	Tropical and Subtropical Fruit, Inedible Peel Group: Papaya	
Target Nematode	Target Plant Pathogens	Application Rate
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 1 -	Root and Tuber Vegetables Group: Potato	
Target Nematode	Target Plant Pathogens	Application Rate
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 1 -	Root and Tuber Vegetables Group: (Exclude Potato, Sugar Beet, and Artichoke) Arracacha, Arrowroot, Bean and True Yam, Beets, Bitter and Sweet Cassava, Black and Spanish Salsify, Carrot, Celeriac (Celery Root), Chayote (Root), Chicory, Chufa, Dasheen (Taro), Edible Burdock, Edible Cana, Ginger, Ginseng, Green Beet, Horseradish, Leren, Oriental Radish, Parsnip, Radish, Rutabaga, Salsify (Oyster plant), Skirret, Sweet Potato, Tanier (Cocoyam), Turmeric, Turnip, Turnip-rooted Chervil, and Turnip-rooted Parsley, and other root and tuber vegetable crops	
Target Nematode	Target Plant Pathogens	Application Rate
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 6 -	Legume Vegetables (Succulent or Dried): Soybeans	
Target Nematode	Target Plant Pathogens	Application Rate
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 13-07 -	Berry and Small Fruit Crop Group: Strawberry	
Target Nematode	Target Plant Pathogens	Application Rate
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 1 -	Root and Tuber Vegetables Group: Sugarbeet	
Target Nematode	Target Plant Pathogens	Application Rate
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*]	1.43 - 5.71 oz/A
<i>Heterodera</i> [*]	<i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	

*Not registered for use in California

Note to Reviewer: Language within brackets is optional label language. Kansas 3 WP Reg. No. 2375-7 Master Label – dated June 27, 2025

Crop -	Sugarcane	
Target Nematode	Target Plant Pathogens	Application Rate
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop -	Tobacco	
Target Nematode	Target Plant Pathogens	Application Rate
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*] <i>Globodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

Crop Group 14 -	Tree Nuts Group: Almond, Beech nut, Brazil nut, Butternut, Cashew, Chinquapin, Coconut, Filbert (hazelnut), Hickory nut, Macadamia nut, Pecan, Walnut, and other tree nuts crops	
Target Nematode	Target Plant Pathogens	Application Rate
<i>Meloidogyne</i> [*] <i>Pratylenchus</i> [*] <i>Heterodera</i> [*]	<i>Rhizoctonia</i> spp.[*] <i>Fusarium</i> spp.[*] <i>Pythium</i> spp.[*] <i>Collectotrichum</i> spp.[*] <i>Verticillium</i> spp.[*]	1.43 - 5.71 oz/A

*Not registered for use in California

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store in a cool, dry place under low humidity. Avoid direct sunlight and avoid excess heat. Keep out of reach of children and animals. Store in original container only. Carefully open container. After partial use, replace lid of container and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal. In case of spill, avoid contact, isolate area, and keep out animals and unprotected persons. Confine spills. Call CHEMTREC (Transportation and Spills) at 1-800-424-9300. To confine spill, dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. If a leaky container must be contained within another, mark the outer container to identify the contents.

Pesticide Disposal

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling

Nonrefillable Plastic Containers (Containers equal to or less than 5 gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer container for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill or by incineration. If burned, stay out of smoke.

Nonrefillable Plastic Containers (Containers greater than 5 gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Follow Pesticide Disposal instructions for rinsate disposal. Repeat this procedure two more times. Then offer container for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill or by incineration. If burned, stay out of smoke.

Conditions of Sale and Limitation of Warranty and Liability

Notice: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of Chr. Hansen or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and, to the extent consistent with applicable law, Buyer and User agree to hold Chr. Hansen and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the compositional description on the label and is reasonably fit for the purposes stated in the Directions for Use when used in accordance with the directions under normal conditions of use.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, CHR. HANSEN MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or Chr. Hansen, and, to the extent consistent with applicable law, Buyer assumes the risk of any such use.

To the extent consistent with applicable law, Chr. Hansen or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CHR. HANSEN AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CHR. HANSEN OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

Chr. Hansen – Trademark of Chr. Hansen Corporation
© 2020 Chr. Hansen Corporation All rights reserved.