

70342-E

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)
CAUTION**

Do not use on humans. Avoid contamination of feed and foodstuffs. Avoid inhalation of spray mist. Avoid contact with skin or eyes. Avoid contact with concentrate. Vapor or spray may be absorbed through the skin. Harmful if swallowed. Wash hands thoroughly after using and before eating or smoking.
Do not use on cats. Use only as directed on the label. Harmful if swallowed. Do not apply to nursing mothers or on puppies under three months of age. Do not use on sick or convalescing animals.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SPRAYING OR DIPPING

Always use the strength recommended on Dilution Table. Add GAMMA R-X directly to cool water to obtain the proper dilution, wt./wt. Complete and thorough coverage of the animal is essential. Treat as necessary, but not more often than once every two weeks for fleas, ticks, and lice.

MIXING - GAMMA R-X mixes easily with cool water, hard or soft. Only slight agitation is required to assure complete suspension of active ingredients. When mixed according to Dilution Table, GAMMA R-X WILL NOT CLOG A SPRAYER.

Dilution Table:
Gamma R-X 1cc 5cc 20cc 8 fl. oz. 1 gal.
Water 7 fl. oz. 1 qt. 1 gal. 12 gal. 200 gal.

Water animals prior to being dipped to avoid animal's drinking of the emulsion. Discard the emulsion when it becomes too fouled for use, or depleted of active ingredient.

TICKS and FLEAS - Mix GAMMA R-X according to dilution table. This emulsion will give control of ticks and fleas present, and is protection against reinfestation.

LICE - Mix GAMMA R-X according to dilution table. Dipping or spraying will kill LICE on the animal. Make a second application in 14 to 18 days to assure control of new LICE hatching from eggs.

SARCOPTIC MANGE - Mix GAMMA R-X according to dilution table, and repeat application at weekly intervals for two or more treatments as necessary.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not contaminate water by disposal of used dip solution. Do not reuse empty container; wrap with newspaper and put in trash collection.

EPA Reg. No. 2382-4
EPA EST. 2382-1N-1

Lot No.

8511 3-1-1 1/2 GALLON



DIP OR SPRAY FOR DOGS

Active Ingredients:
Gamma Isomer of Benzene Hexachloride (from Lindane) 8.82
Aromatic petroleum distillate solvent 86.30
Inert Ingredients 4.88
100.00

GAMMA R-X is a concentrate formulated for spontaneity of Emulsification with a minimum of agitation, forming a stable, long lasting emulsion.
Sold Only By Graduate Veterinarians


KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

- If swallowed - Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. **DO NOT INDUCE VOMITING** vomiting may cause aspiration pneumonia.
- If inhaled - Remove victim to fresh air. Apply artificial respiration if indicated.
- If on skin - Remove contaminated clothing and wash affected areas with soap and water.
- If in eyes - Flush eyes with plenty of water. Get medical attention if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS



1 Gal.