7033-A

SPRAYING OR DIPPING: Always use the strength recommended on Dilution Table. Add F-T-SEVIN directly to cool water to obtain the proper dilution, wt./wt. Complete and thorough coverage of the animal is essential.

Dilution Table:11481F-T-SEVIN\_\_\_\_\_teaspoonful tablespoonful fl. oz.fl. oz.fl. oz.gal.Water\_\_\_\_\_1 pt.3 pts.2½ gal.5 gal.80 gal.

Mix according to dilution table and treat as necessary, but not more often than once every two weeks.

TICKS, FLEAS AND LICE: Mix F-T-SEVIN according to the dilution table. This emulsion will give control of ticks, fleas and lice present and protection against reinfestation. Repeat application as necessary, but not more often than once every two weeks.

Mixing-F-T-SEVIN mixes easily with cool water,

hard or soft. Only slight agitation is required to assure complete suspension of active ingredients.

F-T-SEVIN is recommended as a spray for dog houses, runs and kennels for the control of such ectoparasites as ticks, fleas and lice at a dilution twice the strength used for dipping.

CAUTION TO HUMANS: Avoid contact with concentrate. May be absorbed through the skin. Do not leave in eyes or on skin. Avoid breathing spray mist. Harmful if swallowed. In case of contact wash immediately with soap and water. If taken internally, give 1 tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Call physician immediately. Avoid contamination of foodstuffs.

Note to Physicians: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

CAUTION: Do not reuse container. Destroy when empty.

## WARNING: DO NOT USE ON CATS OR NURSING PUPPIES. SOLD ONLY BY GRADUATE VETERINARIANS

Lot No.

E P A Reg. No. 2382-7 E P A EST. 2382-IN 1



## 

## **CAUTION** KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL FOR ADDITIONAL CAUTIONS

## 1 Gal.

CA DN CHEMICALS, INC. New Castle, Indiana 47362 Made in U.S.A.

F · ··- · 2382-7

61171118; 13