

DIRECTIONS: Dust liberally over animal and rub thoroughly into skin. Be sure to include the ears, legs, feet and between the toes. Comb out dead ticks a few hours after treatment. Repeat application at one to three week intervals if necessary. To control ticks or fleas around animal quarters, remove debris and use at the rate of 4 ounces per 25 sq. ft.

CAUTION: Avoid treatment of puppies or kittens less than 4 weeks old.

CAUTION: Carbaryl is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to other cholinesterase inhibiting drugs, pesticides, or chemicals. Consult veterinarian at the first sign of any adverse reaction. Atropine is the antidote.

CAUTION TO HUMANS: Avoid getting the concentrate on the skin, breathing powder or contamination of foodstuffs. Harmful if swallowed or inhaled. Wash thoroughly after using. In case of contact with eyes, flush with water. If irritation persists, get medical attention. Store in a cool, dry place.

U.S.D.A. Reg. No. 2382-18



FOR USE ON DOGS OR CATS

Especially designed for the control of resistant brown Dog Ticks, Fleas, and Lice. Pleasantly scented to aid in the elimination of "animal odors."

ACTIVE INGREDIENTS: Carbaryl
(1-Naphthyl N-Methylcarbamate) 5.0%
INERT INGREDIENTS 95.0%

CAUTION
KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL for ADDITIONAL CAUTIONS

NET WT. 4 OZ.

2/19/71