

Reg # 2382-95

PM -13

193

Generic pouch label

Flea and Tick Collar For Cats with EPA*

Kills fleas for 5 months

Kills ticks for 5 months

For neck size up to 14"
(optional)

ACTIVE INGREDIENT:

Diazinon: [0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate] 11.0%

INERT INGREDIENTS* 89.0%

TOTAL 100.0%

*Includes 5% unsaturated fatty acids as a grooming aid

KEEP OUT OF REACH OF CHILDREN


CAUTION

See back panel for additional precautionary
statements and directions.

Virbac, Inc.

EPA Reg. No 2382-95 EPA Est. 15577-FR-1
Contains One Flea and Tick Collar Net Wt 0.74 oz. (21g)
Made in France

9504



193

Back of carton

PREVENTEF® Flea and Tick Collar for Cats with EFA

Adjustable - will accommodate up to a 14" neck size

PRECAUTIONARY STATEMENTS

(optional)

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar. May cause contact sensitization following repeated contact with skin in susceptible individuals. Avoid repeated contact with skin. If sensitization reactions result, contact a physician.

Note to Physicians/Veterinarians: Diazinon is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered but only in conjunction with atropine.

Do not use other cholinesterase-inhibiting drugs, pesticides or chemicals on cat while collar is worn. Some cats and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing animals. Do not use on Persian cats.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this collar in a manner inconsistent with its labeling.

Place PREVENTEF® collar around the cat's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and cat's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days. Fleas and ticks present in the animal's environment that may reappear on the cat will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall, however, it is suggested the collar be removed before bathing.

This PREVENTEF® collar contains Essential Fatty Acids necessary for healthy skin and hair. Unsaturated fatty acids released from the collar are rapidly absorbed to help maintain a lustrous, manageable coat.

DISPOSAL

Do not reuse empty pouch or box. Wrap pouch or box and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 2382-95

EPA Est. 15577-FR-1

Manufactured for Virbac, Inc.
P.O. Box 162059, Fort Worth, Texas 76161
Made in France

95/04

Front of carton

PREVENTEF®

Flea and Tick Collar For Cats with EFA*

*Slow release coat conditioner helps enhance texture, manageability and luster

Kills fleas for 5 months

Kills ticks for 5 months

For neck size up to 14"
(optional)

ACTIVE INGREDIENT:

Diazinon [0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate] 11.0%

INERT INGREDIENTS* 89.0%

TOTAL 100.0%

*Includes 5% unsaturated fatty acids as a grooming aid

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

virbac, Inc.

Contains One Flea and Tick Collar Net Wt 0.74 oz (21 g)

95/04

