

PP 13 2382-97

10/1

BLISTER PACKAGE

FRONT PANEL (CARDBOARD BACKING)

**PROTECTION™ Plus 150 Flea and Tick Collar for Dogs
with EPA***

* Slow release coat conditioner helps enhance texture,
manageability and luster.

Kills fleas for 5 months
Kills ticks for up to 5 months

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-
methyl-4-pyrimidinyl) phosphorothioate]...11.0%

INERT INGREDIENTS*.....89.0%

TOTAL.....100.0%

* Includes 5% unsaturated fatty acids as a grooming aid.

KEEP OUT OF REACH OF CHILDREN

**CAUTION: See back panel for additional precautionary statements
and directions.**


VIRBAC

**Contains 1 Flea and Tick Collar
Net Wt. 1.47 oz. (41.5 g)**

ACCEPTED

SEP 9 1991

U.S. DEPARTMENT OF AGRICULTURE
Animal and Plant Health Inspection Service
Washington, D.C. 20250
for the
Director
File No. 2382-97



BLISTER PACKAGE

BACK PANEL (CARDBOARD BACKING)

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing dogs. Do not use on puppies less than 5 weeks of age.

Adjustable - fits all size dogs, up to 28" neck.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labelling.

Place Protectiontm Plus 150 collar around dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against tick will occur within a few days and will continue killing and controlling ticks for up to 150 days; except for unusually large breeds, on which collar aids tick control in the head and neck area for up to 2 months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

This Protectiontm Plus 150 collar contains Essential Fatty Acids necessary for healthy skin and hair. Unsaturated fatty acids released from the collar are rapidly absorbed to help maintain a lustrous, manageable coat.

DISPOSAL

Do not reuse empty blister. After fitting collar, wrap blister and card backing with excess collar and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 2382-97

EPA Est. 15577-FR-1

Manufactured for Virbac, Inc.
2507 Gravel Drive
Fort Worth, Texas 76118
Made in France

24.04.11

FOIL POUCH/CARTON PACKAGE

FRONT PANEL (CARTON)

**PROTECTION™ Plus 150 Flea and Tick Collar for Dogs
with EFA***

* Slow release coat conditioner helps enhance texture,
manageability and luster.

Kills fleas for 5 months
Kills ticks for up to 5 months

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-
methyl-4-pyrimidinyl) phosphorothioate]..11.0%

INERT INGREDIENTS89.0%


* Includes 5% unsaturated fatty acids as a grooming aid.

KEEP OUT OF REACH OF CHILDREN

**CAUTION: See back panel for additional precautionary statements
and directions**

VIRBAC

**Contains 1 Flea and Tick Collar
Net Wt. 1.47 oz. (41.5 g)**



FOIL POUCH/CARTON PACKAGE

BACK PANEL (CARTON)

PROTECTIONtm Plus 150 Flea and Tick Collar for Dogs with EPA.
Adjustable - fits all size dogs, up to 28" neck.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing dogs. Do not use on puppies less than 5 weeks of age.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labelling.

Place Protectiontm Plus 150 collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely to that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days, except for unusually large breeds, on which the collar aids in tick control in the head and neck area for up to 2 months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

6 of 7

This Protection[™] plus 150 collar contains Essential Fatty Acids necessary for healthy skin and hair. Unsaturated fatty acids released from the collar are rapidly absorbed to help maintain a lustrous, manageable coat.

DISPOSAL

Do not reuse empty pouch or box. After fitting collar, wrap pouch and box with excess collar and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 2382-97

EPA Est. 15577-FR-1

Manufactured for Virbac, Inc.
2507 Gravel Drive
Fort Worth, Texas 76118
Made in France

FOIL POUCH/CARTON PACKAGE

POUCH LABEL

**PROTECTIONtm Plus 150 Flea and Tick Collar for Dogs
with EPA**

Do not open until ready to use

CAUTION

KEEP OUT OF THE REACH OF CHILDREN

**SEE BACK PANEL OF OUTER CARTON FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**

ACTIVE INGREDIENT:

**Diazinon [O,O-diethyl O-(2-isopropyl-6-
methyl-4-pyrimidinyl) phosphorothioate]..11.0%**

INERT INGREDIENTS89.0%

TOTAL.....100.0%

Includes 5% unsaturated fatty acids as a grooming aid

EPA Reg. No. 2382-97

EPA Est. 15577-FR-1

**Manufactured for Virbac, Inc.
2507 Gravel Drive
Fort Worth, TX 76118
Made in France**