

PH 13 2382-98

1077

BLISTER PACKAGE

FRONT PANEL (CARDBOARD BACKING)

PROTECTION™ 150 Flea and Tick Collar for Cats

**Kills fleas for 5 months
Kills ticks for 5 months**

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]..15.0%

INERT INGREDIENTS.....85.0%

TOTAL.....100.0%


KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

For neck size up to 12".

VIRBAC

**Contains 1 Flea and Tick Collar
Net Wt. 0.43 oz. (12.2 g)**



ACCEPTED

SEP 9 1991

2382-98

BLISTER PACKAGE

BACK PANEL (CARDBOARD BACKING)

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: Diazinon is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Do not use on cats less than one year old. Do not use other pesticides on cat while collar is worn. Some cats and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing animals. Do not use on Persian cats.
Adjustable - will accommodate up to a 12" neck size.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labelling.

Place Protection[™] 150 collar around the cat's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and cat's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days. Fleas and ticks present in the animal's environment that may reappear on the cat will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

DISPOSAL

Do not reuse empty blister. Wrap blister and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.


20911

EPA Reg. No. 2382-98

EPA Est. 15577-FR-1

Manufactured for Virbac, Inc.
Fort Worth, Texas 76181
Made in France

Page 2



4047

FOIL POUCH/CARTON PACKAGE

FRONT PANEL (CART N1)

PROTECTION™ 150 Flea and Tick Collar for Cats

**Kills fleas for 5 months
Kills ticks for 5 months**

For neck size up to 12".

ACTIVE INGREDIENT:


Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate].....	15.0%
INERT INGREDIENTS.....	85.0%
TOTAL.....	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

VIRBAC

**Contains 1 Flea and Tick Collar
Net Wt. 0.43 oz. (12.2 g)**



FOIL POUCH/CARTON PACKAGE

BACK PANEL (CARTON)

PROTECTION™ 150 Flea and Tick Collar for Cats
Adjustable - will accommodate up to a 12" neck size.

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: Diazinon is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Do not use on cats less than one year old. Do not use other pesticides on cat while collar is worn. Some cats and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing animals. Do not use on Persian cats.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labelling.

Place Protection™ 150 collar around the cat's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and cat's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days. Fleas and ticks present in the animal's environment that may reappear on the cat will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.


DISPOSAL

Do not reuse empty pouch or box. Wrap pouch or box and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 2382-98

EPA Est. 15577-FR-1

Manufactured for Virbac, Inc.
Fort Worth, TX 76181
Made in France



7 of 7

FOIL POUCH/CARTON PACKAGE

POUCH LABEL

PROTECTION™ 150 Flea and Tick Collar for Cats

Do not open until ready to use

CAUTION

KEEP OUT OF REACH OF CHILDREN

**SEE BACK PANEL OF OUTER CARTON FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate].....	15.0%
INERT INGREDIENTS.....	85.0%
TOTAL.....	100.0%

EPA Reg. No. 2382-98

EPA Est. 15577-FR-1

**Manufactured for Virbac, Inc.
Fort Worth, Texas 76181
Made in France**

