FOIL POUCH/CARTON PACKAGE

POUCH LABEL

PROTECTIONtm Flea and Tick Collar for Cats

Do not open until ready to use

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL OF OUTER CARTON FOR ADDITIONAL PRECAUTIONARY STATEMENTS

ACTIVE INGREDIENT:	
Diazinon [0,0-diethyl 0-(2-isopropyl-6-	
methyl-4-pyrimidinyl) phosphorothioate]15.()ફ
INERT INGREDIENTS85.0	
TOTAL	

EPA Reg. No. 2382-98

EPA Est. 15577-FR-1

Manufactured for Virbac, Inc. Fort Worth, Texas 76181 Made in France

ACCEPTED

FEB 25 1993

Under the Federal Insecticide. Fungicide, and Rodenticlica Acc. as amended, for the poeticide registered under EPA Reg. No. 2232-98

FOIL POUCH/CARTON PACKAGE

FRONT PANEL (CARTON)

PROTECTIONtm 150 Flea and Tick Collar for Cats

Kills fleas for 5 months Kills ticks for 5 months

STEP 3 logo

For neck size up to 12*

ACTIVE INGREDIENT:

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

VIRBAC

Contains 1 Flea and Tick Collar Net Wt. 0.43 oz. (12.2 g)

----<u>-</u>

* **

FOIL POUCH/CARTON PACKAGE

BACK PANEL (CARTON)

Zone Control: To effectively control fleas, you should treat the areas where fleas live with the Flea Stop^T 4-Step Program which consists of, Step 1 -Shampoo/Dip the Pet; Step 2 -Treat the Home and Yard; Step 3 -Flea and Tick Collar on the Pet; Step 4 - Maintenance Sprays and Powders. Ask your pet care professional for specific recommendations on how to best control fleas for the comfort and well-being of your pet and family.

PROTECTIONtm 150 Flea and Tick Collar for Cats Adjustable - will accomodate up to a 12" neck size.

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESIC ANIMALS

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar. May cause contact sensitization following repeated contact with skin in susceptible individuals. Avoid repeated contact with skin. If sensitization reactions result contact a physician.

Note to Physicians/Veterinarians: Diazinon is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Do not use other cholinesterase inhibit_ng pesticides, drugs or chemicals on cat while collar is worn. Some cats and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing animals. Do not use on Persian cats.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labelling.

Place Protectiontm 150 collar around the cat's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and cat's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly.



Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days. Fleas and ticks present in the animal's environment that may reappear on the cat will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

DISPOSAL

Do not reuse empty pouch or box. Wrap pouch and box and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 2382-98

EPA Est. 15577-FR-1

Manufactured for Virbac, Inc. Fort Worth, TX 76181 Made in France