

BLISTER PACKAGE

FRONT PANEL (CARDBOARD BACKING)

PROTECTION™ 150 Flea and Tick Collar for Large Dogs

**Kills fleas for 5 months
Kills ticks up to 5 months**

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]..15.0%

INERT INGREDIENTS.....85.0%

TOTAL.....100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

For neck size up to 28".

VIRBAC

**Contains 1 Flea and Tick Collar
Net Wt. 0.90 oz. (25.5 g)**

ACCEPTED

SEP 9 1991

**U.S. DEPARTMENT OF AGRICULTURE
OFFICE OF THE SECRETARY
WASHINGTON, D.C. 20250
PHONE 2382-99**

BLISTER PACKAGE

BACK PANEL (CARDBOARD BACKING)

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: Diazinon is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing animals.
Adjustable - fits large dogs, 28" neck or smaller.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labelling.

Place Protectiontm 150 collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days. except for unusually large breeds on which collar aids in tick control in the head and neck area for up to 2 months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

DISPOSAL

Do not reuse empty blister. Turn up blister and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

FOIL POUCH/CARTON PACKAGE

FRONT PANEL (CARTON)

PROTECTIONtm 150 Flea and Tick Collar for Large Dogs

**Kills fleas for 5 months
Kills ticks for up to 5 months**

For neck size up to 28".

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate].....15.0%

INERT INGREDIENTS.....85.0%


TOTAL.....100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

VIRBAC

**Contains 1 Flea and Tick Collar
Net Wt. 0.90 oz. (25.5 g)**



FOIL POUCH/CARTON PACKAGE

BACK PANEL (CARTON)

PROTECTIONtm 150 Flea and Tick Collar for Large Dogs
Adjustable - fits large dogs, 28" neck or smaller.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: Diazinon is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing animals.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labelling.

Place Protectiontm 150 collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days, except for unusually large breeds on which collar aids in tick control in the head and neck area for up to 2 months. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.


DISPOSAL

Do not reuse empty pouch or box. Wrap pouch or box and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 2382-99

EPA Est. 15577-FR-1

Manufactured for Virbac, Inc.
Fort Worth, TX 76181
Made in France



7/2/18

FOIL POUCH/CARTON PACKAGE

POUCH LABEL

PROTECTION™ 150 Flea and Tick Collar for Large Dogs

Do not open until ready to use

CAUTION

KEEP OUT OF REACH OF CHILDREN

**SEE BACK PANEL OF OUTER CARTON FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-	
methyl-4-pyrimidinyl) phosphorothioate].....	15.0%
INERT INGREDIENTS.....	85.0%
TOTAL.....	100.0%

EPA Reg. No. 2382-99

EPA Est. 15577-FR-1

**Manufactured for Virbac, Inc.
Fort Worth, Texas 76181
Made in France**

Faint, illegible text or markings in the bottom right corner of the page.

PH 13 2520 90

BLISTER PACKAGE

FRONT PANEL (CARDBOARD BACKING)

PROTECTIONtm 150 Flea and Tick Collar for Dogs

**Kills fleas for 5 months
Kills ticks for 5 months**

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]	..15.0%
INERT INGREDIENTS.....	85.0%
TOTAL.....	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

For neck size up to 22".

VIRBAC

**Contains 1 Flea and Tick Collar
Net Wt. 0.85 oz. (24.0 g)**

ACCEPTED

SEP 9 1991

2382-99

9.24.74

BLISTER PACKAGE

BACK PANEL (CARDBOARD BACKING)

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: Diazinon is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing animals.
Adjustable - fits dogs, 22" neck or smaller.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labelling.

Place Protectiontm 150 collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.


DISPOSAL

Do not reuse empty blister. Wrap blister and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 2382-99

EPA Est. 15577-FR-1

Manufactured for Virbac, Inc.
Fort Worth, Texas 76181
Made in France



FOIL POUCH/CARTON PACKAGE

FRONT PANEL (CARTON)

PROTECTION™ 150 Flea and Tick Collar for Dogs

**Kills fleas for 5 months
Kills ticks for 5 months**

For neck size up to 22".

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate].....15.0%

INERT INGREDIENTS.....85.0%


TOTAL.....100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

VIRBAC

**Contains 1 Flea and Tick Collar
Net Wt. 0.84 oz. (24.7 g)**



12518

FOIL POUCH/CARTON PACKAGE

BACK PANEL (CARTON)

PROTECTION™ 150 Flea and Tick Collar for Dogs
Adjustable - fits dogs, 22" neck or smaller.

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.

Note to Physicians/Veterinarians: Diazinon is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing animals.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labelling.

Place Protection™ 150 collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.


DISPOSAL

Do not reuse empty pouch or box. Wrap pouch or box and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 2382-99

EPA Est. 15577-FR-1

Manufactured for Virbac, Inc.
Fort Worth, TX 76181
Made in France



FOIL POUCH/CARTON PACKAGE

POUCH LABEL

PROTECTION™ 150 Flea and Tick Collar for Dogs

Do not open until ready to use

CAUTION

KEEP OUT OF REACH OF CHILDREN

**SEE BACK PANEL OF OUTER CARTON FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**


ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate].....15.0%
INERT INGREDIENTS.....85.0%
TOTAL.....100.0%

EPA Reg. No. 2382-99

EPA Est. 15577-FR-1

**Manufactured for Virbac, Inc.
Fort Worth, Texas 76181
Made in France**



BLISTER PACKAGE

FRONT PANEL (CARTON)

PROTECTIONtm 150 Flea and Tick Collar for Puppies

Kills fleas for 5 months
Kills ticks for 5 months

For puppies over 5 weeks - neck size up to 12".

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate].....	15.0%
INERT INGREDIENTS.....	85.0%
TOTAL.....	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.


VIRBAC

Contains 1 Flea and Tick Collar
Net Wt. 0.46 oz. (13.0 g)

ACCEPTED

SEP 9 1991

2-382-99



FOIL POUCH/CARTON PACKAGE

BACK PANEL (CARTON)

PROTECTION™ 150 Flea and Tick Collar for Puppies
Adjustable - will accomodate up to a 12" neck size.

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar.


Note to Physicians/Veterinarians: Diazinon is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Do not use other pesticides on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing puppies. Do not use on puppies less than 5 weeks of age.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labelling.

Place Protection™ 150 collar around the puppy's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 150 days). Ticks are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days and will continue killing and controlling ticks for up to 150 days. Fleas and ticks present in the animal's environment that may reappear on the puppy will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.




DISPOSAL

Do not reuse empty pouch or box. Wrap pouch or box and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 2382-99

EPA Est. 15577-FR-1

Manufactured for Virbac, Inc.
Fort Worth, TX 76181
Made in France



FOIL POUCH/CARTON PACKAGE

POUCH LABEL

PROTECTIONtm 150 Flea and Tick Collar for Puppies

Do not open until ready to use

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE BACK PANEL OF OUTER CARTON FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

ACTIVE INGREDIENT:

Diazinon [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate].....	15.0%
INERT INGREDIENTS.....	85.0%
TOTAL.....	100.0%

EPA Reg. No. 2382-99

EPA Est. 15577-FR-1

Manufactured for Virbac, Inc.
Fort Worth, Texas 76181
Made in France

