

Reg # 2382-102

PM - 13

191

BEST AVAILABLE COPY

virbac

DuoCide® Dip

**Residual Insecticidal Flea and Tick Dip
for Dogs and Cats for 14 day Control and
Repellency by Synergised Permethrin**

ACTIVE INGREDIENTS:

Permethrin [(1R)-3-Phenoxyphenyl] methyl (±) cis-trans-3-(2,2-dichloroethenyl) 2,2- dimethylcyclopropanecarboxylate].....	3.20%
Related reaction products.....	0.28%
n-Octyl Bicycloheptene Dicarboximide.....	15.00%
INERT INGREDIENTS.....	81.25%
TOTAL.....	100.00%

*cis-trans isomer ratio: Min. 35% (±) cis
Max. 65% (±) trans

WARNING

KEEP OUT OF REACH OF CHILDREN.

**See back panel for additional precautionary
statements.**

Net Wt. 4 fl. oz. (114 mL)

Made in U.S.A. by: virbac®, inc.
P.O. Box 162059, Fort Worth, Texas 76161

PRECAUTIONARY STATEMENTS WARNING
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
 Causes skin irritation. Harmful if swallowed. Do not get in eyes, on skin or on clothing. Do not contaminate feedstuffs with this concentrate or dip. Do not use on animals under 3 months of age or on sick, or on old or debilitated pets.

ENVIRONMENTAL HAZARDS
 This pesticide is toxic to fish and other aquatic animals. Do not contaminate any body of water with this concentrate or dip or by cleaning of equipment.

STATEMENT OF PRACTICAL TREATMENT
IF SWALLOWED: Call a physician Or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. **DO NOT INDUCE VOMITING.** Vomiting may cause aspiration pneumonia.
IF ON SKIN: Remove contaminated clothing and wash affected areas with soap and warm water.
IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.
IF INHALED: Remove victim to fresh air and call a physician if effects occur. Apply artificial respiration if necessary.

DIRECTIONS FOR USE
 It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
 To kill fleas, ticks and lice. Dilute 2 fl. oz (2 tablespoons) with one gallon of water. Dip animal making sure hair is wet to the skin. The area around the ears should be sponged thoroughly. Dipping should not exceed 30 seconds. Avoid contact of dip with eyes and mouth of animal. If a volume of dip less than one gallon is required, e.g. to apply to smaller animals or by sponging onto the animal rather than immersion dipping, maintain a dilution rate of one oz per half gallon or 1/4 oz. per quart of water. Allow pet to dry without rubbing or toweling. Repeat application every 2 weeks or as necessary.

STORAGE AND DISPOSAL DIRECTIONS
STORAGE: Store in closed container in a cool dry area
DISPOSAL: Do not reuse empty container. Wrap container in newspaper and put in trash.
 EPA Reg. No. 2382-102 EPA Est. No. 2382-IN-1
 Lot No. on bottom of container 83/11

ACCEPTED

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 2382-102