

Reg # 2382-104

PM-19

181

PREVENTIC
TICK COLLAR For Dogs
 Neck Size Up to 25

Kills Ticks for 3 Months

ACTIVE INGREDIENT:
 Amitraz N-(2,4-dimethyl-phenyl)-N-[[2,4-dimethyl-phenyl]imino]meth]-N-methylmethanimidamide 9.0%

INERT INGREDIENTS 91.0%

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION: See back panel for additional precautionary statements and directions

virbac, Inc.

CONTENTS ONE COLLAR (NET WT 0.97 OZ (27.5g))

Tick Collar for Dogs
 Adjustable - Fits dogs, 25" neck or smaller

PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION: This collar is intended for use as an acaricide generator and is not to be taken internally by man or animal. Not intended for use on humans. Do not allow children to handle or play with this collar. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Some humans may develop skin irritation through contact with collared pet. Remove collar, shampoo pet and consult physician if irritation persists. Wash thoroughly with soap and water after handling.

Some dogs may be sensitive to this collar. Remove at first sign of irritation or adverse reaction and shampoo dog thoroughly to remove residue. Do not use on sick or convalescing dogs. **DO NOT APPLY TO CATS.** There is a risk of collar breakage and ingestion if collar is worn by unattended dogs that play by mouthing each other in the neck area. If collar is missing from a dog, search for, retrieve and recover the entire collar. Immediately consult a veterinarian if dog is depressed.

Note to Veterinarian: If a dog ingests a Preventic collar and exhibits intoxication (primary sign is severe depression), induce vomiting, administer iron only laxative, activated charcoal and enema. Provide supportive therapy until collar fragments are voided. If dog cannot be roused, Yohimbine, 0.1 mg/kg by intravenous injection is antidotal. Repeat as needed.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this collar in a manner inconsistent with its labeling.

Flare the Preventic® collar around the dog's neck, buckle and adjust for proper fit. Cut off and dispose of excess length. Do not allow dog to play with cut off excess. The collar must be worn tightly enough to contact skin and so that dog cannot remove it. Activity against ticks starts within 24 hours and continues as long as the collar is worn (up to 90 days). Ticks present in the dog's environment that may reappear on the dog will not attach and will be killed in less than 24 hours. Replace collar when effectiveness diminishes. The effectiveness of the collar is not diminished by normal wetting such as rainfall; however, it is suggested that the collar be removed before bathing. The active ingredient is not an insecticide. Therefore, use other means for control of fleas.

DISPOSAL

Do not reuse empty pouch, box or collar. Wrap in newspaper and put in trash collection.

EPA Reg. No. 2382-104 EPA Est. 15577-FR-1
 Manufactured for virbac®, inc.
 P. O. Box 162053, Fort Worth, Texas 76161
 Made in France

LOT

2410

ACCEPTED

JAN 04 1995

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 2382-104.