

FOIL POUCH/CARTON PACKAGE

FRONT PANEL (CARTON)

PROTECTION™ 300
Flea and Tick Collar for Dogs

Kills fleas for 10 months
Kills Brown Dog Ticks for 7 months

For neck size up to 28"

ACTIVE INGREDIENT:

Diazinon [0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate].....15.0%

INERT INGREDIENTS.....85.0%

TOTAL.....100.0%


KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

VIRBAC

CONTAINS 1 FLEA AND TICK COLLAR
NET WT. 1.4 OZ. (42 g)

ACCEPTED
SEP 25 1992
Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act, this product is registered for use as follows:
EPA Reg. No. 2382-105



FOIL POUCH/CARTON PACKAGE

BACK PANEL (CARTON)

PROTECTIONtm 300 Flea and Tick Collar for Dogs
Adjustable - fits large dogs, 28" neck or smaller.

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar. May cause contact sensitization following repeated contact with skin in susceptible individuals. Avoid repeated contact with skin. If sensitization reactions result contact a physician.

Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotal only if signs of cholinesterase inhibition are present.

Do not use other cholinesterase inhibiting pesticides, drugs or chemicals on dog while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing animals. Do not use on puppies under one year of age. Do not use more than one collar per animal.

DIRECTIONS FOR USE

It is a violation of Federal law to use this collar in a manner inconsistent with its labelling.

Place Protectiontm 300 collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Activity against fleas starts within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 300 days). Brown Dog Ticks (*Rhipicephalus sanguineus*) are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days. This collar will continue killing and controlling ticks for up to 210 days. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal wetting such as rainfall; however, it is suggested the collar be removed before bathing.

BEST AVAILABLE COPY

