

For Dogs

RELEASE™ Flea and Tick Collar for Dogs

Adjustable - fits dogs, 20" neck or smaller.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: This collar is intended for use as an insecticide/growth regulator and is not to be taken internally by man or animal. Not intended for use on humans. The insecticide in this collar is a cholinesterase inhibitor. Do not allow children to handle or play with this collar. May cause dizziness or drowsiness following repeated contact with skin in susceptible individuals. Avoid repeated contact with skin. If sensitization reactions occur, consult a physician. Note to Physicians/Veterinarians: This product contains a cholinesterase inhibitor. Atropine sulfate and 2-PAM are antidotes only if signs of cholinesterase inhibition are present.

Do not use other cholinesterase-inhibiting pesticides, drugs or chemicals on dogs while collar is worn. Some dogs and humans may be sensitive to this collar. Remove at first sign of irritation or adverse reaction. Do not use on sick or convalescing animals. Do not use on puppies under one year of age. Do not use more than one collar per animal.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this collar in a manner inconsistent with its labeling. Place the RELEASE™ collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of collar as length. The collar must be worn loosely so that two fingers can be placed between collar and dog's neck. If collar is worn too tightly, it may produce neck irritation. Active against flea eggs within 24 hours, reaches maximum effectiveness in 3-4 days and continues as long as the collar is worn (up to 250 days). Brown Dog Ticks (*Rhipicephalus sanguineus*) are relatively harder to kill and die more slowly. Activity against ticks will occur within a few days. This collar will continue killing and controlling ticks for up to 210 days. Fleas and ticks present in the animal's environment that may reappear on the dog will be killed and controlled by the action of the collar. Replace collar when effectiveness diminishes. Collar is not affected by normal washing such as rainfall; however, it is suggested the collar be removed before bathing.

STORAGE AND DISPOSAL: Storage: Store in a cool dry area.

Disposal: Do not reuse empty pouch or box. Wrap pouch or box and excess collar length and put in trash collection. When discarding used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 2382-105

EPA Est. 1987-FR-1

Manufactured for Virbac, Inc.
2307 Grand Drive
Fort Worth, Texas 76116
Made in Mexico

LOT:

0210

RELEASE™
Flea and Tick Collar for Dogs

Kills fleas for 10 months
Kills Brown Dog Ticks for 7 months
For dogs up to 25 lbs

ACTIVE INGREDIENT

KEEP OUT OF REACH OF CHILDREN
CAUTION: See back panel for additional product safety information.

virbac, Inc.

ACCEPTED
JAN 19 1993

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 2382-105

BEST AVAILABLE COPY