

virbac[®]

PREVENDER Residual Flea and Tick Foam For Dogs

Active Ingredient:
Permethrin [(3-Phenoxyphenyl) methyl
(±) cis-trans-3-(2,2-dichloroethenyl) 2,2-
dimethylcyclopropanecarboxylate] .0.677%
Inert Ingredients 99.323%

*cis-trans isomer ratio: Max. 55% (±) cis
Min. 45% (±) trans

**KEEP OUT OF REACH OF CHILDREN
WARNING
SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**

EPA Reg. No. 2382-118
EPA Est. No. 2382-IN-1

Net Weight: 9 fl. oz. (223 gm.)

Made in USA for
Virbac, Inc.
P.O. Box 162059, Fort Worth, Texas 76161

KILLS FLEAS AND AMERICAN AND BROWN DOG TICKS FOR 2 WEEKS

Precautionary Statements and Domestic Animals Hazards to Humans
WARNING

Hazards to Humans: Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Due to irritating nature may be harmful if swallowed. Wash hands thoroughly with soap and water after handling.

Hazards to Domestic Animals: Avoid contact with eyes. Do not use on puppies under 6 weeks-old. In the event of reactions, foam residue may be removed by washing the dog with a non-insecticidal shampoo. If reaction persists consult a veterinarian.

Statement of Practical Treatment: If in eyes: Flush with plenty of water for 15 minutes. Call a physician. If swallowed: Drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol.

Physical or Chemical Hazards: Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Do not store below 32°F. Exposure to temperature above 130°F may cause bursting.

Directions for Use: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **Shake well** and depress nozzle to dispense. Foam can be applied directly onto the pet or into the palm of your hand and then applied to the pet. For direct application, apply one-second bursts of Prevender Flea Foam as hair is backcombed. Apply over the entire animal until damp, massaging or brushing the hair until the foam vanishes. Allow pet to dry. Do not apply from can directly to face, rather apply foam by hand around ears and under chin. This contents of this can (applied at a rate of 7gm foam/kg body weight) will provide approximately 4 treatments or 2 months protection against fleas and ticks for an 18 lb dog. Prevender Flea Foam may be reapplied every 2 weeks or as necessary when the pet is reinfested.

Storage and Disposal: **Storage:** Store in a cool area away from heat or open flame. Do not store below 32°F. **Disposal:** Do not reuse empty container. Replace cap, wrap container in several layers of newspaper and discard in trash. Do not puncture or incinerate. L of no 95/01

FEB 1 1995

2382-118

BEST AVAILABLE COPY