Thomas A. Miller, DVM, Ph.D. Virbac Inc. P.O. Box 162059 Fort Worth, Texas 76161

Subject: Revised Label and CSF For:

Flea Ovisterilant Collar for Cats #1

EPA Registration No. 2382-135

Application dated February 26, 1997

Dear Dr. Miller:

The amendments to the registration for the product cited above under the Federal Insecticide, Fungicide, And Rodenticide Act, as amended are acceptable. The amended labeling is acceptable subject to the comment listed below:

Delete the phrase, is not toxic, from the sentence, The active ingredient in this collar, Nylar, xxx adult parasites.

A copy of the stamped draft label is enclosed for your records. Submit five copies of the corrected labeling.

The Confidential Statements Of Formula dated February 26, 1997 are acceptable and have been placed into the file for this product.

The additional brand name, KnockOut Flea Collar for Cats and Kittens, is acceptable.

If you have any questions or comments please contact Joseph Tavano at (703) 305-6411.

Sincerely,

RPK

Richard P. Keigwin, Jr.
Product Manager 10
Insecticide & Rodenticide Branch
Registration Division (7505C)

Flea Ovisterilant Collar For Cats #1 IGR Flea Collar for Cats and Kittens

[Sterilizes fleas for 13 months] [Sterilizes (and kills) flea eggs for 13 months]

ACTIVE INGREDIENT: Pyriproxyfen: 2-[1-methyl-2-(phenoxy-phenoxy) ethyoxy] 0.5% pyridine..... INERT INGREDIENTS.

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for additional precautionary statements and directions.

> virbac* Contains One Flea Ovisterlant Collar Net Wt. 0.42 oz. (12g)

> > 97/02

BEST AVAILABLE COPY

ACCEPTED with CONTRENTS In EPA Letter Dated

MAR 1 7 1997 : topoglicide, Under the T The Jack Fragist? el enimero registered under Erich g. No. 2382-135

Carton, back

READ ENTIRE LABEL BEFORE USING USE ONLY ON CATS

PRECAUTIONARY STATEMENTS CAUTION
HAZARDS TO HUMANS: This collar is intended for use only on pet animals. Do not allow readwards to Hosakes. This color is transcen for use only on per animals. Do not allow obtained to handle or play with this color. Wash hands with soep and water after handling. HAZARDS TO ANIMALS: Do not use on nursing kittens. Consult a veterinarian before using on disblittated, aged, medicated, pregnant or nursing animats. Sensitivities may occur after using any pesticide product on pets. If signs of sensitivity occur, remove coller and battle your pet with mild soep or a non-pesticidal shampoo and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

DIRECTIONS FOR USE it is a violation of Federal Law to use this collar in a manner inconsistent with its labelling. Place collar around the cat's neck, adjust for proper fit and buckle in place. Out off and dispose of excess langth. The collar must be worn lightly enough to contact skin.

If coller is worn too lightly, it may produce neck inflation.
This coller is dealgned for long term prevention and control of flee problems by preventing the build up of flee populations through its adult flee and flee egg sterifizing actions. Ideally, as a preventive measure and for meadrum effect, this coller should be applied to the cat before the The season starts. Its files sterifizing activity, that sterifizes files season starts. Its files sterifizing activity, that sterifizes files stready on the cat, starts immediately, reaches 100% effectiveness within 2 days and confinues at 100% effectiveness as long as the collar is worn (at least for 370 days). Filess stready present in the cats anytoment, that may response no files cut, will be immediately sterifized before they start to lay eggs so that only sterifie dead eggs will be taid. The potential for development of new files in the cat's environment, that would normally reinfest the cut, will thus be prevented for a full year. It is recommended therefore that this collar should be replaced once a year, perhaps on a aignificant date such as the pet's birthday or on the first of January.

The active ingredient in this collar, Nylar*, is not an insecticide or acericide, is not toxic and will noticil adult parasites. Therefore, if the cat or its environment is heavily infested with flees when the collar is first applied, the collar's flee and flee egg sterilizing benefit will not be apparent for several weeks until all pre-sudsing environmental flee life cycle stages have developed and either found a host or died. In these circumstances it is necessary to use other means (e.g., etments, applied to both the pet and its environment) to kill existing fies stages on the pet and in its environment. Tick infestation must always be controlled by products labelled for ticks. The collar and its active ingredient in the cat's haircost are not affected by normal weiting or by beithing. Replace collar as needed.

DISPOSAL Do not reuse empty pouch or box.
When pouch or box and put in trash collection. When discarding out-off excess collar or used ooliar, wrap in newspaper and place in trash collection. EPA Reg. No. 2382-135 Virbec, inc.

EPA Est, 2382-TX-1

55 Virbec, inc. P.O. Box 152059 Fort Worth, Texas 76161

Telephone 817-831-5030 blyler*, is a registered trademark of McLaughtin Gormley King Company.

Generic pouch label

Flea Ovisterilant Collar For Cats #1 IGR Flea Collar for Cats and Kittens

[Sterilizes fleas for 13 months] [Sterilizes (and kills) flea eggs for 13 months]

ACTIVE INGREDIENT:	
Pyriproxyfen: 2-[1-methyl-2-(phenoxy-phenoxy) ethyoxy]	
pyridine	0.59
NERT INGREDIENTS	99.59

KEEP OUT OF REACH OF CHILDREN CAUTION

See carton for additional precautionary statements and directions.

EPA Reg. No. 2382-135

EPA Est. 15577-FR-1

Contains One Fiea Ovisterliant Collar Net Wt. 0.42 oz. (12g)

Made in France

97/02