

generic pouch label Enlarged to 150%

AMITRAZ TICK COLLAR FOR DOGS

Neck Size up to 25'

Kills Ticks for 2 Months

ACTIVE INGREDIENT:

Amitraz: N'-(2,4-dimethylphenyl)-N-[[2,4-dimethyl-phenyl)imino]methyl]-N-methylmethanimidamide	6.0%
INERT INGREDIENTS	94.0%

CAUTION

KEEP OUT OF REACH OF CHILDREN

See back panel of carton for additional precautionary statements and directions.

EPA Reg. No. 2382-139

EPA Est. 2382-FR-1


virbac, Inc.

CONTENTS: ONE COLLAR

NET WT. 0.97 OZ (27.5g)

9702

ACCEPTED
 MAR 11 1997
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 2382-139



2 7 2

back of carton Enlarged to 150%

**PRECAUTIONARY STATEMENTS READ ENTIRE LABEL BEFORE EACH USE
USE ONLY ON DOGS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION**

HAZARDS TO HUMANS: This collar is intended for use as an acaricide generator and is not to be taken internally by man or animal. Not intended for use on humans. Do not allow children to handle or play with this collar. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Some humans may develop skin irritation through contact with collared pet. Remove collar, shampoo pet and consult physician if irritation persists. Wash thoroughly with soap and water after handling.

HAZARDS TO PETS: DO NOT APPLY TO CATS. Do not apply to puppies under 12 weeks of age. Consult a veterinarian before using on debilitated, aged, medicated, pregnant or nursing animals. There is a risk of collar breakage and ingestion if collar is worn by unattended dogs that play by mouthing each other in the neck area. If collar is missing from a dog, search for, retrieve and recover the entire collar. Immediately consult a veterinarian if dog is depressed. Sensitivities may occur after using any pesticide product on pets. If signs of sensitivity occur bathe pet with a non-pesticidal shampoo or mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

Note to Veterinarian: If a dog ingests a Preventic collar and exhibits intoxication (primary sign is severe depression), induce vomiting, administer non-oily laxative, activated charcoal and enema. Provide supportive therapy until collar fragments are voided. If dog cannot be roused, Yohimbine, 0.1 mg/kg by intravenous injection is antidotal. Repeat as needed.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this collar in a manner inconsistent with its labeling. Place the collar around the dog's neck, buckle and adjust for proper fit. Cut off and dispose of excess length. Do not allow dog to play with cut-off excess. The collar must be worn tightly enough to contact skin and so that dog cannot remove it. Activity against ticks starts within a few days and continues as long as the collar is worn (up to 60 days). Ticks present in the dog's environment that may reappear on the dog will not attach and will be killed in 2-3 days. Replace collar when effectiveness diminishes. The effectiveness of the collar is not diminished by normal wetting such as rainfall, however, it is suggested that the collar be removed before bathing. The active ingredient is not an insecticide. Therefore, use other means for control of fleas.

DISPOSAL: Do not reuse empty pouch, box or collar. Wrap in newspaper and put in trash collection.

EPA Reg. No. 2382-139

EPA Est. 2382-FR-1

Manufactured for virbac®, Inc.
P. O. Box 162059, Fort Worth, Texas 76161
Made in France

Lot number stamped on pouch

97/02

