

Preventic PLUS⁺

TICK AND FLEA IGR COLLAR FOR DOGS

READ ENTIRE LABEL DIRECTIONS BEFORE EACH USE. USE ONLY ON DOGS. DO NOT USE ON CATS.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION:

HAZARDS TO HUMANS: Not intended for use on humans. Do not allow children to handle or play with this collar. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Some humans may develop skin irritation through contact with collared pet. Remove collar, shampoo pet and consult physician if irritation persists. Wash thoroughly with soap and water after handling.

HAZARDS TO PETS: Do not use on puppies less than 12 weeks old. Consult a veterinarian before using on debilitated, aged, medicated, pregnant or nursing animals. Do not treat collared dogs with other monoamine oxidase inhibitors, tricyclic antidepressants, selective serotonin reuptake inhibitors or pressor agents. Remove at first sign of irritation or adverse reaction, bathe dog with a non-pesticidal shampoo and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. There is a risk of collar breakage and ingestion if collar is worn by unattended dogs that play by mouthing each other in the neck area. If collar is missing from a dog, search for, retrieve and recover the entire collar. Immediately consult a veterinarian if dog is depressed.

Note to Veterinarian: If a dog ingests a Preventic Plus collar and exhibits intoxication (primary sign is severe depression), induce vomiting, administer non-oily laxative, activated charcoal and emera. Provide supportive therapy until collar fragments are voided. If dog cannot be roused, Yohimbine, 0.1mg/kg by intravenous injection is antidotal. Repeat as needed. Temporary and reversible hyperglycemia was observed when the product was tested at 5 times the recommended dose.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this collar in a manner inconsistent with its labeling. The two active ingredients in this collar have different and complementary functions. The IGR (synthetic flea hormone), pyriproxyfen (Nylar®), sterilizes all fleas that jump on the pet (even new fleas before they start laying eggs) and kills all flea eggs. Pyriproxyfen is continuously released from the collar, spreads over the pet in the natural oils of the haircoat and skin and adheres lightly to hair and skin to provide more than 3 months of continuous flea ovicidal efficacy. This breaks the flea's life cycle by preventing the development of new fleas in the environment that would normally reinfect the pet. This residual activity of pyriproxyfen starts immediately, reaches 100% effectiveness within 3-4 days and continues at 100% effectiveness as long as the collar is worn. The second active, amitraz, also spreads over the pet in the oil of its skin and haircoat where its activity starts within 24 hours to detach and kill existing ticks on the pet. Amitraz continues to prevent attachment and feeding and kills new ticks that may climb onto the pet for up to 90 days. The collar and its active ingredients are not affected by normal wetting such as rainfall, however it is suggested the collar be removed before bathing. This collar should be replaced when new ticks are seen to persist on the pet. For optimal efficacy for tick control and flea prevention you should replace the collar every 3 months.

DIRECTIONS FOR USE: Place the collar around the dog's neck, adjust for proper fit and buckle in place. Cut off and dispose of excess length. Do not allow dogs to play with cut-off excess. The collar must be worn tightly enough to contact skin and so the dog cannot remove it.

DISPOSAL AND DISPOSAL STORAGE

Store in a cool, dry place. **DISPOSAL:** Do not reuse empty pouch or box. Wrap pouch and put in trash collection. When discarding cut off end and used collar, wrap in newspaper and place in trash collection.

EPA Reg. No. 2382-170

EPA Est. 2382-TX-1

Manufactured for

virbac

P.O. Box 162059
Fort Worth, Texas 76161
Telephone (817) 831-5030

Made in France

Nylar® is a registered trademark of
McLaughlin Gormley King Company

99/05

Preventic PLUS⁺ TICK AND FLEA IGR COLLAR FOR DOGS

ANTIPARASITIC



Preventic[®] PLUS

Kills ticks on dogs for 3 Months
Nylar[®] IGR sterilizes fleas and
kills flea eggs on dogs for 3 months

ACTIVE INGREDIENTS:

Pyriproxyfen	0.5%
Amitraz	9.0%
INERT INGREDIENTS	90.5%

KEEP OUT OF REACH OF CHILDREN

CAUTION: See back panel for additional precautionary statements and directions.

EPA Reg. No. 2382-170

EPA Est. 2382-TX-1

Contains One Amitraz-Pyriproxyfen Collar
Net Wt. 1.5 oz. (42g)

Made in France

virbac

Preventic[®] PLUS⁺ TICK AND FLEA IGR COLLAR FOR DOGS

2382-170

11/16/2000

Page 131

ANTONE 661 CV

PREVENTIC PLUS COLLAR (U.S.) 10/15/2000

ACCEPTED

NOV 16 2000

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 2382-170

FOR VETERINARY USE ONLY