


Hopkins

2393-108

PM 23 1063

7½% CAPTAN DUST



ACCEPTED
Per Self Cont
EPA 2 1 1987
2393-108

ACTIVE INGREDIENTS

Captan	7.33%
Related Derivatives	0.17%
INERT INGREDIENTS	92.50%
Total	100.00%

N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

KEEP OUT OF REACH OF CHILDREN
DANGER PELIGRO

PRECAUCION AL USUARIO: Si usted no lee Ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Call a physician.
If swallowed: Call a physician or Poison Control Center immediately. Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available, drink large quantities of water. Avoid alcohol. Do not give anything by mouth to an unconscious person.
If inhaled: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.
If on skin: Wash thoroughly with soap and water.

See Side Panel for Additional Precautionary Statements
EPA Reg. No. 2393-108
EPA Est. No. 10543-MO-1

STORE IN A COOL DRY PLACE.
PROTECT FROM EXCESSIVE HEAT.



SOLD BY
Hopkins
Agricultural Chemical Co.

2001 7332 - MADISON, WISCONSIN 53701
U.S.A.

NET WEIGHT 50 POUNDS

2063

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

DANGER

Causes irreversible eye damage. Avoid if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes.
Avoid contact with skin and clothing.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Avoid consumption of food and feed stuffs.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bays, marshes, and ponds). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.
In the event that treated seeds are spilled and cannot be salvaged, completely cover them with soil.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not allow persons to enter treated areas within 4 days following application unless a long-sleeved shirt and long pants or a cover-all that covers all parts of the body except the head, hands and feet and chemically resistant gloves are worn.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral precautions must be given to workers engaged in agricultural hand labor who are expected to be in a treated area or in an area about to be treated with this product. These precautions are:
DANGER: area treated with poison on (date of application).

Do not enter treated areas within 4 days following application unless a long-sleeved shirt and long pants or a cover-all that covers all parts of the body except the head, hands and feet and chemically resistant gloves are worn.

For accidental exposure, see Statement of Practical Treatment.
Oral precautions must be given if there is reason to believe that written precautions cannot be understood by workers. When oral precautions are given, precautions shall be given in a language customarily understood by workers.

Wear goggles or face shield when handling.
Wash thoroughly with soap and water after handling.
Remove contaminated clothing and separately launder clothing before reuse.

Mixers/loaders when mixing and loading this product must wear a long sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron.

Fieldworkers and harvesters must wear chemical resistant gloves. Leather or fabric gloves are not acceptable.

A dust mask must be worn when mixing and loading this product.

During application only, from a tractor with a completely enclosed cab or from an aircraft with an enclosed cockpit a long sleeved shirt and long legged pants must be worn. Chemically resistant gloves must be available in the cab or cockpit and worn while exiting.

AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with captan must be destroyed according to state and local regulations.

CROP DIRECTIONS:

APPLE: Apple Scab, Black Rot (Frogeye Leaf Spot) Black Pox, Bitter Rot, Botryosphera, Brooks Fruit Spot, Flyspeck—25 to 50 lbs. per acre (9.2 ozs. to 1 lb. 2.4 ozs./1,000 sq. ft.). Make first application at delayed dormant period. Repeat applications for each infection period or at weekly intervals as long as necessary.

APRICOT: Brown Rot, Jacket Rot—50 to 60 lbs. per acre (1 lb. 2.4 ozs. to 1 lb. 6 ozs./1,000 sq. ft.). Apply at red bud stage and repeat at 75 percent petal fall.

CHERRY: Brown Rot, Leaf Spot—25 to 50 lbs. per acre (9.2 ozs. to 1 lb. 2.4 ozs./1,000 sq. ft.). Apply at popcorn, blossom and petal fall stages; continue at 10 to 14 day intervals.

PEACH: Brown Rot, Scab—25 to 50 lbs. per acre (9.2 ozs. to 1 lb. 2.4 ozs./1,000 sq. ft.). Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make pre-harvest application where necessary.

PEAR: Pear Scab (except Pacific Northwest)—25 to 40 lbs. per acre (9.2 ozs. to 14.7 ozs./1,000 sq. ft.). Make first applications during early finger and petal fall stages. Repeat applications for each infection period or at weekly intervals as long as necessary. Russeting may be produced on Bosc pears. Do not use on D'Anjou pears.

STRAWBERRIES: Botrytis Rot—30 to 40 lbs. per acre (11.0 ozs. to 14.7 ozs./1,000 sq. ft.). Make application when new growth starts in spring before fruit starts to form. Repeat applications weekly. Severe infection conditions—Continue through harvest period, treating immediately after each picking.

CUCURBITS (Cantaloupe, Cucumber): Angular Leaf Spot, Anthracnose, Downy Mildew — Start application when runners begin to form or at the first sign of disease. Apply at rate of 20 to 30 lbs. per acre (7.35 ozs. to 11.0 ozs./1,000 sq. ft.) on young plants, increasing to 40 lbs. per acre (14.7 ozs./1,000 sq. ft.) on mature plants.

POTATO SEED PIECE TREATMENT: To protect against seed piece breakdown and rot or decay from Brown Eye Disease, Damp-off, Seed Rot, Verticillium Wilt — Use at rate of 10 ozs. per bushel cut seed pieces (16 ozs. or 1 lb. to 100 lbs. cut seed pieces). Apply in convenient container or by duster attachment over belt. Coat each seed piece evenly with a light film of dust. Treat pieces within 6 hours after cutting. If planting is delayed, store treated seed pieces in open crates for two days before bagging. **DO NOT USE TREATED SEED FOR FOOD OR FEED PURPOSES.**

TOMATOES: Anthracnose, Early and Late Blight, Gray Leaf Spot, Septoria Leaf Spot — Apply 30 to 40 lbs. per acre (11.0 ozs. to 14.7 ozs./1,000 sq. ft.) at weekly intervals on young plants; 40 to 50 lbs. per acre (14.7 ozs. to 1 lb. 2.4 ozs./1,000 sq. ft.) on mature plants.

DO NOT USE FOR ANY PURPOSE NOT RECOMMENDED ON THIS LABEL, SINCE INJURIOUS EFFECTS MAY RESULT.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a cool, dry place in original container. Protect from excessive heat. Once opened, do not store bag for prolonged period.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE OF WARRANTY

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instruction.