

# Hopkins

## PROLIN<sup>®</sup> PELLETS

Price  
**\$1.79**

### RAT and MOUSE KILLER A NEW, IMPROVED Warfarin Rodenticide In Pellet Form

ACTIVE INGREDIENTS:

Warfarin 3-( <del>acetonyl</del> benzyl)-4-hydroxycoumarin . . . . .	0.025
N-(2-quinoxaliny) sulfanilamide (sulfaquinoxaline) . . . . .	0.025

INERT INGREDIENTS . . . . .	99.950
Total . . . . .	100.000

\* Prolin is the registered trademark of the Wisconsin Alumni Research Foundation for a rodenticide.  
USDA Reg. No. 2393-145

### CAUTION

KEEP OUT OF REACH OF CHILDREN,  
DOMESTIC ANIMALS AND PETS

See Side Panels for Complete Caution Statement,  
Use Directions and Notice

NET WEIGHT 2 LBS.

PROLIN contains Warfarin, an anticoagulant chemical that reduces the clotting ability of the blood and, upon repeated feedings, causes fatal hemorrhage in rats and mice. It also contains an antibacterial agent, sulfaquinoxaline, which inhibits bacteria that produces Vitamin K, an antidote for Warfarin. It thereby increases effectiveness against hard-to-kill rats and mice and gives better and more thorough control.

NOTICE: When used as directed Seller warrants this product to be effective for the purposes stated hereon. Seller makes no other warranties of any kind, express or implied, concerning the use of this product. Buyer assumes all risk of use or handling whether in accordance with directions or not.

24-8A7

### USE DIRECTIONS

#### CAUTION

Place baits in locations not accessible to children, pets, and domestic animals, or in tamper-proof bait boxes. If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K, combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin. In case of ingestion, call a physician.

RATS: Bait generously. Place 4 to 6 ounces of Hopkins Prolin Pellets in dry container next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. Replace stale, damp or moldy baits with fresh bait.  
MICE: Place tablespoonful amounts of Hopkins Prolin Pellets in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 15 days. Mice are "scatterers" so make plenty of bait placements to assure continuous feeding.

HOPKINS AGRICULTURAL CHEMICAL CO  
Madison, Wisconsin