

pot 14

2393-149

1/3

BEST AVAILABLE COPY

Hopkins

PROLIN[®] ANTICOAGULANT CONCENTRATE



STATEMENT OF PRACTICAL TREATMENT

If on Skin: Wash thoroughly with soap and water.

If Swallowed: Call a physician or Poison Control Center immediately. Induce vomiting by drinking 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If in Eyes: Flush with water.

NOTE TO PHYSICIAN: If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intramuscular and oral administration of Vitamin K₁ combined with blood transfusions are indicated, as in the case of hemorrhage caused by overdose of bishydroxycoumarin. Repeat as necessary based on monitoring of prothrombin times.

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS & DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. Exposure during pregnancy should be avoided. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. When mixing baits, wear dust respirator and rubber gloves. If on skin, wash with soap and water.

ENVIRONMENTAL HAZARD Keep Out of Lakes, Streams or Ponds.

NOTICE OF WARRANTY

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks of use and/or handling of this material when such use and/or handling is contrary to label instructions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MIXING DIRECTIONS

Mix 1 part Hopkins Prolin Concentrate with 19 parts (by weight) of a suitable bait material such as freshly ground whole yellow corn, rolled oats, or other dry grain materials attractive to rats and mice. If very dusty, add up to 5% of a bland edible oil such as mineral or corn oil. One pound makes 20 lbs. of mixed bait.

As an example of ingredients to be used: mix thoroughly 1 part of this product with 17 parts whole coarse ground wheat and 2 parts of powdered sugar.

Bait materials resembling human food must be altered in form by crushing, balling, or pelleting so that they are not readily recognizable as food.

Directions continued on right panel

ACTIVE INGREDIENT:

Warfarin [3-(alpha-Acetylbenzyl)-4-hydroxycoumarin]	0.5%
INERT INGREDIENTS:	99.5%
TOTAL	100.0%

Contains 0.5 Sulfiquinoxaline
Trademark of the Wisconsin Alumni Research Foundation for a rodenticide

**KEEP OUT OF REACH OF CHILDREN,
DOMESTIC ANIMALS AND PETS**

CAUTION

See Side Panels for Additional Cautions and Use Directions

EPA Reg. No. 2393-149
EPA Est. No. 2393-B-31-2393-WI-1*
2393-WI-2-33509-MO-1*
Superscript refers to first letter of lot number



Hopkins

agricultural chemicals

NET WEIGHT

50 POUNDS
(22.7 Kilograms)

Dire
USE
rats, home
tured
dump
rabb
Treat
bait
child
not
poss
come
SEL
Dete
most
these
ings,
ceak
local
obse
Rem
RAT
ment
bait
MIC
spoc
plac
point
unit
days
RAT
spoi
all d
To
and
tion
tinu
perm
need
Do
sto
PE
no
mu
ap
of
Ac
Co
ba
to
int
ba
all
CO

Directions (continued from left panel)

USE RESTRICTIONS: For control of Norway rats, Roof rats and House mice in and around homes, buildings and similar man-made structures, in transportation vehicles, or sites such as dumps, sewers, river and pond banks, feedlots, rabbitry fur farms, and game bird operations.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals and wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS — Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 1/4 to 1/2 ounce (1-2 level table-spoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS & MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

BEST AVAILABLE COPY

77-100-1001-72

ACCEPTED

NOV 18 1982

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No. **2343-149**