


Hopkins

PROLIN

ANTICOAGULANT CONCENTRATE



1.00% (w/v) Sodium Warfarin
 2.00% (w/v) Sodium Coumatetralil Hydrochloride
 1.00% (w/v) Sodium Phosphotungstate Hexahydrate
 1.00% (w/v) Sodium Phosphate Dibasic Heptahydrate
 1.00% (w/v) Sodium Chloride
 1.00% (w/v) Sodium Citrate
 1.00% (w/v) Sodium Bicarbonate
 1.00% (w/v) Sodium Hydroxide
 1.00% (w/v) Sodium Metaphosphate
 1.00% (w/v) Sodium Sulfate
 1.00% (w/v) Sodium Borate
 1.00% (w/v) Sodium Acetate
 1.00% (w/v) Sodium Citrate
 1.00% (w/v) Sodium Bicarbonate
 1.00% (w/v) Sodium Hydroxide
 1.00% (w/v) Sodium Metaphosphate
 1.00% (w/v) Sodium Sulfate
 1.00% (w/v) Sodium Borate
 1.00% (w/v) Sodium Acetate

CAUTION

KEEP OUT OF REACH OF CHILDREN, DOMESTIC ANIMALS & PETS

Treated bait should be stored in locations not accessible to children, pets, wildlife and poultry. If tampered with, it should be destroyed by humans, domestic animals, wildlife or pets. The bait may reduce the clotting ability of the blood and cause bleeding internally and externally. Treatment of Vitamin K combined with blood transfusions are indicated as to the extent of hemorrhage caused by overdoses of bishydroxycoumarin.

This rodenticide contains Vitamin K, an anticoagulant, which is fatal to rats. The anticoagulant, upon repeated feedings, causes fatal hemorrhage in rats and mice. It also contains a substance which inhibits the production of Vitamin K in the intestine of the rat.

Do not use in poultry, swine, dogs, cats, horses, or other animals. Destroy any bait remaining in the container immediately after use.

For further information, contact your local distributor or Hopkins Agricultural Chemical Co., Madison, Wisconsin 53701.

DIRECTIONS FOR USE

Use 1 quart (32 oz) of concentrate per 100 lbs. of bait. Mix with 100 lbs. of bait. The bait should be stored in a cool, dry place. Do not use in poultry, swine, dogs, cats, horses, or other animals. Destroy any bait remaining in the container immediately after use.

FOR CONTROLLING RATS: Apply 1 quart (32 oz) of concentrate per 100 lbs. of bait. Mix with 100 lbs. of bait. The bait should be stored in a cool, dry place. Do not use in poultry, swine, dogs, cats, horses, or other animals. Destroy any bait remaining in the container immediately after use.

FOR CONTROLLING MICE: Apply 1 quart (32 oz) of concentrate per 100 lbs. of bait. Mix with 100 lbs. of bait. The bait should be stored in a cool, dry place. Do not use in poultry, swine, dogs, cats, horses, or other animals. Destroy any bait remaining in the container immediately after use.

NOTICE OF WARRANTY

Hopkins Agricultural Chemical Co. warrants that the product is as described on the label. Buyer assumes all other risks and is not liable for any loss or damage resulting from the use of the product in any manner not intended by the manufacturer.

NET WEIGHT

POUNDS

HOPKINS AGRICULTURAL CHEMICAL CO.

Madison, Wisconsin 53701

Hopkins

FOR CONTROLLING RATS

For information on the use of this product, please refer to the side panels. This product is a rodenticide and should be used in accordance with the directions on the label. It is not to be used on food or feed. It is not to be used in areas where it may be consumed by humans or animals. It is not to be used in areas where it may be consumed by birds or other animals. It is not to be used in areas where it may be consumed by fish or other aquatic animals. It is not to be used in areas where it may be consumed by insects or other arthropods. It is not to be used in areas where it may be consumed by plants or other organisms. It is not to be used in areas where it may be consumed by soil organisms. It is not to be used in areas where it may be consumed by microorganisms. It is not to be used in areas where it may be consumed by any other organism.

FOR CONTROLLING MICE

For information on the use of this product, please refer to the side panels. This product is a rodenticide and should be used in accordance with the directions on the label. It is not to be used on food or feed. It is not to be used in areas where it may be consumed by humans or animals. It is not to be used in areas where it may be consumed by birds or other animals. It is not to be used in areas where it may be consumed by fish or other aquatic animals. It is not to be used in areas where it may be consumed by insects or other arthropods. It is not to be used in areas where it may be consumed by plants or other organisms. It is not to be used in areas where it may be consumed by soil organisms. It is not to be used in areas where it may be consumed by microorganisms. It is not to be used in areas where it may be consumed by any other organism.

PROLIN[®] REDI-MIX RAT AND MOUSE KILLER

A New, Improved
WARFARIN Rodenticide in Meal Form
READY-TO-USE BAIT

ACTIVE INGREDIENTS:

Warfarin, [3-(α -acetylbenzyl)-4-hydroxycoumarin] 0.025%
N¹-(2-quinoxaliny) sulfanilamide (sulfquinoxaline) 0.025%

INERT INGREDIENTS: 99.950%

Total 100.000%

[®] Prolin is a registered trademark of the Wisconsin Alumni Research Foundation for a rodenticide.

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Side Panels for Complete Caution Statement, Use Directions and Notice. **1 POUND**

HOPKINS AGRICULTURAL CHEMICAL CO.

Madison, Wisconsin

FOR CONTROLLING RATS

Bait generously. Place 4 to 6 ounces of Hopkins Prolin Redi-Mix Rat and Mouse Killer in dry container next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped. Replace stale damp or moldy baits with fresh bait.

FOR CONTROLLING MICE

Place tablespoonful amounts of Hopkins Prolin Redi-Mix Rat and Mouse Killer in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 15 days. Mice are "nibblers" so make plenty of bait placements to assure continuous feeding.

Hacco 10M 967

USDA Reg. No. 2393-150

Hopkins

HOPKINS AGRICULTURAL CHEMICAL CO.
Madison, Wisconsin