2393-166

í

## PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K - intramuscularly or orally, as indicated in bishydroxycoumarin overdose. Repeat as necessary based on monitoring of prothrombin times.

### ENVIRONMENTAL HAZARD

Keep Out of Lakes, Streams or Ponds.

EPA Reg. No. 2393-166 EPA Reg. No. 2393-1L-3<sup>4</sup>, 2393-WI-1<sup>3</sup>, 523-IL-1<sup>25</sup>, 2393-WI-2<sup>19</sup>, 33509-MO-1<sup>4</sup> Superacript refers to first letter of lot number,

rm 14

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS. For control of Norway rats, Roof rats and House mice in and around homes, buildings, and similar man-made structures and in transportation vehicles.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Determine areas where rais and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bail from rain or snow. Remove as much food as possible.



### APP! ICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 1/4 to 1/2 ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days

RATS & MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait property. To prevent reinfestation, eliminate food, water, and harborage as much as possible if reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed

# DISPOSÁL

DO NOT REUSE EMPTY CONTAINER WRAP CONTAINER AND PUT IN TRASH COLLECTION

# NOTICE OF WARRANTY

Manufacturer and Seller make no warranty of merchantability, litness for any purpose, or otherwise, express or implied, concerning this product or its use, which extend beyond the statements on this label

CODE 2393.166 1 2 3 4 5 6 7 8 9 (8/82)

