ACCEPTED

DEC 26 1992

Under the Federal Insecticide, Fungicide, and Redenticide Act, os amended, for the pesticide registered under 213-166

Page 1 of 4 Date: September, 1992

HOPKINS RODEX PELLETED BATT-1

Rat and Mouse Killer

Contains Microencapsulated Warfarin

ACTIVE INGREDIENT:

Warfarin {3-(alpha-Acetonylbenzyl)-4-hydroxycoumarin} 0.025%

INERT INGREDIENTS:

99.975%

TOTAL

KEEP OUT OF REACH OF CHILDREN {12 pt¹}
CAUTION {18 pt¹}

SEE RIGHT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 2393-166

EPA Est. No. 61282-WI-1

NEI CONTENTS

MANUFACTURED FOR: HACO, INC. P.O. BOX 7190 MADISON, WI 53707 U.S.A.

¹Assumes a front panel of over 30 sq. in.

RODEX PELLETED BAIT-1
Page 2 of 4

Date: September, 1992

{Left Panel}

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

- 1. Store product not in use in a location out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
- 3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS - For control of Norway rats, roof rats, and house mice in and around homes, industrial buildings, and similar man-made structures. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS - Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 oz. of bait per placement*. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 1/4 to 1/2 oz. (1-2 level tablespoons) of bait at 8- to 12-foot intervals. Larger placements (up to 2 oz.) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

*For sizes less than 1 lb., the following statement will be added: "The amount of bait in this package is unlikely to be sufficient to kill more than one or two rats."

Rodex Pelleted Bait-1 Page 3 of 4 Date: September, 1992

{Right Panel}

APPLICATION DIRECTIONS (Continued from left panel)

RATS/ Replace contaminated or spoiled bait immediately collect and MICE: dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

(For Horsehold Products¹)

STORAGE AND DISPOSAL {12 pt2

Storage: Store only in original contain r, in a dry

place inaccessible to children and pets.

Disposal: Do not reuse empty container. S curely

wrap in newspaper and discard in rash.

{For Non-Household Products³}

STORAGE AND DISPOSAL {12 pt2}

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep product in tightly closed original container. Store in a dry area not accessible to children, pets, domestic animals or wildlife. Do not store near material with a strong odor.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

¹ For sizes 5 lbs. or less.

² Assumes a front panel of over 30 sq. in.

³ For sizes larger than 5 lbs.

Rodex Pelleted Bait-1

Page 4 of 4

Date: September, 1992

PRECAUTIONARY STATEMENTS: HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN - If ingested, administer Vitamin K₁ intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS

This product is toxic to mammals and birds. Do not apply this product to water, swamps, bogs, or potholes.

NOTICE OF WARRANTY

Seller and manufacturer make no warranty of merchantability, fitness for any purpose, express or implied concerning this product or its use, which extend beyond the statements on this label.

Ready Mixed