


Hopkins

FRUIT TREE SPRAY

MULTI-PURPOSE SPRAY FOR
SMALL HOME ORCHARDS



ACTIVE INGREDIENTS:

Captan (N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide).....	10%
Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate).....	5%
5-Methoxychlor, Technical*	5%
INERT INGREDIENTS:	80%
Total	100%

* 5% Methoxychlor Technical is equivalent to 4.4% of 2,2-bis (p-methoxyphenyl) - 1,1,1-trichloroethane and 0.6% of other isomers and related compounds.

CAUTION

KEEP OUT OF REACH OF CHILDREN AND PETS

SEE BACK PANEL FOR USE INSTRUCTIONS AND TERMS OF SALE

Avoid inhalation of dust or spray mist. Avoid contact with skin. Avoid contamination of feed and foodstuffs. Wash thoroughly after using. Do not apply within 7 days of harvest.

This product is toxic to fish and wildlife. Avoid contamination of fish bearing streams by dust, spray or residues that may drift or wash into water.

NET WEIGHT 3 POUNDS

Hopkins Agricultural Chemical Co.
Madison, Wisconsin

Hopkins

FRUIT TREE SPRAY

MULTI-PURPOSE SPRAY FOR
SMALL HOME ORCHARDS

Jan - 5 - 1968

*2393-
170*


ACTIVE INGREDIENTS

Water
Sulfur
Zinc
Copper
Lime
Oil
Soap

CAUTION

KEEP OUT OF REACH OF CHILDREN

Always use protective clothing and equipment when handling this product. Do not breathe vapors or dusts. Avoid contact with skin and eyes. Wash thoroughly after handling. Do not eat, drink, or use tobacco products while using this product. Dispose of contents and container as directed. Do not reuse container.



Hopkins

FRUIT TREE SPRAY

MULTI-PURPOSE SPRAY FOR
SMALL HOME ORCHARDS

ACTIVE INGREDIENTS:

Captan (N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide)	10%
Malathion (O,O-dimethyl dithophosphate of diethyl mercaptosuccinate)	10%
Methoxychlor, Technical*	10%
INERT INGREDIENTS:	<u>70%</u>
TOTAL	100%

* 10% Methoxychlor Technical is equivalent to 8.8% of 2,2-bis (P-methoxyphenyl)-1,1,1-trichloroethane and 1.2% of other isomers and related compounds.

EPA Reg. No. 2393- 273

EPA Est. No. 2393-WI-1

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Left Panel For Use Instructions
and Notice of Warranty

HARMFUL IF SWALLOWED. Avoid inhalation of dust or spray mist. Avoid contact with skin. Avoid contamination of feed and foodstuffs. Wash thoroughly after using. Do not apply within 7 days of harvest.


This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NET WEIGHT

1 POUND

HOPKINS AGRICULTURAL CHEMICAL CO.
Madison, Wisconsin 53701



HOPKINS FRUIT TREE SPRAY

For protection against codling moth, aphid, bud moth, leaf roller, apple maggot and apple scab.

DIRECTIONS FOR USE

Use 5-6 level tablespoons per gallon of water, or for larger amounts, at the equivalent rate of 4-5 lbs. per 50 gallons of water. Spray until the solution drips off the leaves, making sure both upper and lower surfaces of each leaf are covered.

Do not spray during bloom, but spray at **early** petal fall and then every 7-10 days until 2 weeks before harvest.

The period from green tip to bloom is very important for scab control. During this period, spray every seven days, or more often if growth is rapid or if rains occur frequently. (There may be 3-5 applications before bloom, depending upon growth development or weather conditions.) For primary scab infection, use 5-6 level tablespoons per gallon of water, or for larger amounts, at the equivalent rate of 4-5 lbs. per 50 gallons of water. Apply in prebloom, blossom, petal fall, and first cover periods. For secondary scab infections, use 2.0 level tablespoons per gallon water, or for larger amounts, at the equivalent rate of 1.33 lbs. per 50 gallons of water. Apply in cover periods. Whenever an insect infestation occurs, spray at that time. In the event of reported severe outbreak of apple maggot, make an application of Diazinon (1.0 lb. of Diazinon 50W per 100 gals. of water) when recommended by the State Agricultural Experiment Station.

Red Delicious and other sensitive varieties of apple may be injured by early season application.

NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

HOPKINS AGRICULTURAL CHEMICAL CO.
Madison, Wisconsin 53701