

Contains SEVIN[®] Insecticide
For the Contro. of Certain Vegetable Crop Insect Pests

ACTIVE INGREDIENT:
 Carbaryl (1-naphthyl methylcarbamate) 5%

INERT INGREDIENTS: 95%

TOTAL 100%

[®]SEVIN is the registered trade mark of Union Carbide Corporation for carbaryl insecticide.

EPA Reg. No. 2393-174

CAUTION

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Harmful if inhaled or swallowed. Avoid breathing of dust. Do not take internally. Avoid prolonged or repeated contact with skin. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. Avoid contamination of food, feed, water supplies, streams and ponds.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

DIRECTIONS FOR USE

To control the insects on the crop, specified apply Hopkins 5% Carbaryl Dust at the amounts stated below. Apply in a manner to give uniform coverage. Make first application when insects or their damage appear. Repeat as specified or as necessary.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER: Controls flea beetles and harlequin bug, use at the rate of 12-20 lbs. per acre, for cabbage worms, loopers and corn earworm, use 20-40 lbs. per acre.

BEANS Green, Snap or Lima Controls Japanese beetle, green stink bug, cucumber beetle, flea beetles, Japanese beetle, leaf miner, etc. Use at the rate of 10-20 lbs. per acre.

Apply in a manner to give uniform coverage. Make first application when insects or their damage appear. Repeat as specified or as necessary.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER: Controls flea beetles and harlequin bug, use at the rate of 12-20 lbs. per acre; for cabbage worms, loopers and corn earworm, use 20-40 lbs. per acre.

BEANS – Green, Snap or Lima: Controls Mexican bean beetle, bean leaf beetles, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, armyworms, stink bugs and tarnished plant bug, use at the rate of 20-30 lbs. per acre.

CORN – Sweet Corn: Controls corn earworm, European corn borer, Japanese beetle, sap beetles. To control foliar feeding insects, treat entire plant repeating as necessary. To control insects feeding on ears and silks treat at 2-3 day intervals, beginning when silks appear and continue until silks are dry, use at the rate of 20-40 lbs. per acre; (1/2-1 lb. per 1,000 sq. ft. for small gardens). In small gardens for the control of European corn borer apply thoroughly into the whorls when plants are about knee high. Repeat at weekly intervals until tassels emerge. For field application for the control of European corn borer consult your state agricultural experiment station for proper timing of treatments.

Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

POTATO, TOMATO, EGGPLANT, PEPPER: Controls Colorado potato beetle, flea beetles, leafhoppers, use at the rate of 12-20 lbs. per acre; for tomato fruitworm, tomato hornworm and stink bugs, use 20-40 lbs. per acre.

CUCUMBER, MELONS, PUMPKIN AND SQUASH: Controls pickleworm and melonworm, use at the rate of 12-20 lbs. per acre. For cucumber beetles, flea beetles, leafhoppers and squash bugs, use 20 lbs. per acre. NOTE: Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. Do not overtreat.

NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling of this material is contrary to label instructions.

NET WEIGHT

50 POUNDS

HOPKINS AGRICULTURAL CHEMICAL CO.

Madison, Wisconsin 53701