


Hopkins

10% TRITHION[®] PLUS 5% THIRAM GRANULES

An Organic Phosphate A Soil Insecticide



ACTIVE INGREDIENTS:

Carbophenothion S [(p-chlorophenylthio)methyl] O,O-diethyl phosphorodithioate	10.0
Thiram (Tetramethylthiuram disulfide)	5.0

INERT INGREDIENTS:	85.0%
TOTAL	100.0%

© Reg. U.S. Patent Office, Chemical Company, for insecticide use only.
EPA Reg. No. 2393-173

CAUTION

KEEP OUT OF REACH OF CHILDREN

OBSERVE THE FOLLOWING USE PRECAUTIONS -- HAZARDOUS IF SWALLOWED

Avoid breathing dust, contact with skin and eyes. Wear dust respirator and goggles. Wash hands, arms, and face thoroughly with soap and water after handling and before eating or smoking. Do not contaminate feed and foodstuffs. Wash all contaminated clothing with soap and hot water before re-use. **TRITHION** may cause cholinesterase inhibition. **ATROPINE IS ANTIDOTAL.** Keep a supply of Atropine 1/100 grain tablets on hand.

Do not apply or allow to drift to areas occupied by unprotected humans, beneficial animals or onto adjoining food and forage crops.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label. Do not contaminate water by cleaning of equipment or disposal of wastes.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates.

DIRECTIONS FOR USE

ONIONS: To control onion maggot and onion smut from seeds, sets or transplants (dry bulb or green) -- **FURROW TREATMENT** -- Apply in the planting furrow at time of seeding or transplanting using a combination drill or suitable granular application equipment.

MUCK SOIL: Apply 15 to 20 lbs. per acre for planting in 18 inch rows (9-11 oz. per 1,000 linear feet of row).

MINERAL SOIL: Apply 10 to 15 lbs. per acre for planting in 18 inch rows (6-8 oz. per 1,000 linear feet of row).

NOTICE OF WARRANTY

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all use and/or handling of this material when such use and/or handling is contrary to label instructions.

DESTROY EMPTY CONTAINER BURN STAY OUT OF SMOKE

NET WEIGHT

50 POUNDS

Hopkins Agricultural Chemical Co.

Madison, Wisconsin 53701

Hopkins

SPECIAL GRANULES

For Control of Onion Maggots & Onion Smut
Contains 5% *Dasanit and 5% Thiram

Insecticide-Fungicide

FOR USE ONLY BY COMMERCIAL APPLICATORS

ACTIVE INGREDIENTS:

Dasanit (O, O-Diethyl O-lp-(methylsulfinyl) phenyl phosphorothioate)*	5.0%
Thiram (Tetramethylthiuram disulfide)	5.0%
INERT INGREDIENTS:	90.0%
Total	100.0%

*Dasanit Reg. TM in U.S. Pat. Off. Farbenfabriken Bayer A. G., Chemagro Corporation licensee.
U.S. Patent No. 3,042,703; Canadian Patent No. 631,472
USDA Reg. No. 2393-182

DANGER!

KEEP OUT OF REACH OF CHILDREN



POISON



STOP! READ LABEL COMPLETELY BEFORE USE

Poisonous if swallowed, inhaled, or absorbed through the skin. Rapidly absorbed through the skin. Do not get in eyes, on skin or on clothing. Do not breathe dust. Wear protective clothing and cotton or natural rubber gloves. If cotton gloves are used, they must be laundered or discarded after each day's use. Do not wear the same gloves for other work. Rubber gloves should be washed with soap and water after each use. Wash hands, arms and face thoroughly with soap and warm water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Keep all unprotected persons out of the operating area or vicinity. Do not use in or near human dwellings. When using formulation in greenhouses or other enclosed areas, wear a mask or respirator effective for protection against Dasanit.

Do not contaminate feed or food. Do not use on plants other than those recommended. Use only according to label directions. Application at rates above those shown may result in illegal crop residues.

Toxic to fish and wildlife. Do not contaminate streams, lakes, or ponds. Birds feeding on treated areas may be harmed.

SYMPTOMS OF POISONING: A sense of "tightness" in the chest. Sweating. Contracted pupils. Stomach pains. Vomiting and diarrhea.

TREATMENT: In case of poisoning call a physician immediately. Have patient lie down and keep quiet. If swallowed, induce vomiting by giving a tablespoonful of salt in a glass of warm water. Administer milk or water freely and again induce vomiting until vomit fluid is clear. Never give anything by mouth to an unconscious person. Give artificial respiration if breathing has stopped. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash with flowing water for at least 15 minutes.

NOTE TO PHYSICIAN: ANTIDOTE: Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Compound inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system, and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically.

CONTAINER DISPOSAL

DO NOT RE-USE THE CONTAINER - Completely empty the contents and bury the unused chemical at least 18 inches deep in an isolated location away from water supplies.

Burn the empty container completely. Exercise extreme caution and stay well away from the smoke.

Do not use container in any connection with feed, food, and drinking water.

DIRECTIONS FOR USE

The use of Hopkins Special Granules is limited to use on onions in the North Central and Northeastern States only.

Hopkins Special Granules will give effective control of onion maggots (including those resistant to certain chlorinated hydrocarbon insecticides) and onion smut.

ONION (DRY BULB ONLY): To control onion maggot and onion smut, apply Hopkins Special Granules at the rate of 20 pounds per acre with 18 inch row spacing or at the rate of 10 ounces per 1,000 linear feet of row regardless of row spacing. In the event of extremely narrow row spacing, do not exceed the rate of 20 pounds per acre. Distribute the granules evenly in the seed furrow at planting time. **DO NOT APPLY TO GREEN-BUNCH ONIONS.**

TERMS OF SALE

NOTICE: "Seller" as used in this NOTICE and statement of TERMS OF SALE includes manufacturers and distributors of product. Used

MAY-5-1967
UNDER THE
FUNGICIDE AND
FOR ECONOMIC RATIONING
IS UNDER NO. 2393-182